

## Validation

1. HEARD-LIPSMEYER M, et al. Evaluating body composition in infancy and childhood: A comparison between 4C, QMR, DXA, and ADP. *Pediatr Obes.* 2020 Jun;15(6):e12617. doi: 10.1111/ijpo.12617. Epub 2020 Jan 27. PMID: 31986239; PMCID: PMC7323309
2. FORSUM E, et al. An evaluation of the Pea Pod system for assessing body composition of moderately premature infants. *Nutrients.* 2016 Apr 22;8(4):238. doi: 10.3390/nu8040238. PMID: 27110820; PMCID: PMC4848706
3. FIELDS D, et al. Body composition at 6 months of life: Comparison of air displacement plethysmography and dual-energy x-ray absorptiometry. *Obesity (Silver Spring).* 2012 Nov;20(11):2302-6. doi: 10.1038/oby.2012.102. Epub 2012 Apr 23. PMID: 22522885
4. ROGGERO P, et al. Evaluation of air-displacement plethysmography for body composition assessment in preterm infants. *Pediatr Res.* 2012 Sep;72(3):316-20. doi: 10.1038/pr.2012.75. Epub 2012 Jun 5. PMID: 22669294
5. FRONDAS-CHAUTY A, et al. Air displacement plethysmography for determining body composition in neonates: Validation using live piglets. *Pediatr Res.* 2012 Jul;72(1):26-31. doi: 10.1038/pr.2012.35. Epub 2012 Mar 22. PMID: 22441376
6. ANDRES A, et al. Quantitative nuclear magnetic resonance to measure fat mass in infants and children. *Obesity (Silver Spring).* 2011 Oct;19(10):2089-95. doi: 10.1038/oby.2011.215. Epub 2011 Jul 21. PMID: 21779094
7. ELLIS K, et al. Body-composition assessment in infancy: Air-displacement plethysmography compared with a reference 4-compartment model. *Am J Clin Nutr.* 2007 Jan;85(1):90-5. doi: 10.1093/ajcn/85.1.90. PMID: 17209182
8. YAO M, et al. Inter-device reliability of the PEA POD® for percent body fat estimates. <http://www.cosmed.com> (2005)
9. MA G, et al. Validation of a new pediatric air-displacement plethysmograph for assessing Body Composition in Infants. *Am J Clin Nutr.* 2004 Apr;79(4):653-60. doi: 10.1093/ajcn/79.4.653. PMID: 15051611
10. YAO M, et al. Preliminary evaluation of a new pediatric air displacement plethysmograph for body composition assessment in infants. *Acta Diabetol.* 2003 Oct;40 Suppl 1:S55-8. doi: 10.1007/s00592-003-0027-9. PMID: 14618434
11. URLANDO A, et al. A new air displacement plethysmograph for the measurement of body composition in infants. *Pediatr Res.* 2003 Mar;53(3):486-92. doi: 10.1203/01.PDR.0000049669.74793.E3. PMID: 12595599
12. SAINZ R, et al. Evaluation of a new pediatric air-displacement plethysmograph for body-composition assessment by means of chemical analysis of bovine tissue phantoms. *Am J Clin Nutr.* 2003 Feb;77(2):364-70. doi: 10.1093/ajcn/77.2.364. PMID: 12540395

## Validation of Other Techniques Using the PEA POD

1. VAN BEIJSTERVELDT I, et al. Skinfold-based-equations to assess longitudinal body composition in children from birth to age 5 years. *Clin Nutr.* 2023 Jul;42(7):1213-1218. doi: 10.1016/j.clnu.2023.04.024. Epub 2023 Apr 29. PMID: 37225558.
2. RODRIGUEZ-CANO A, et al. Development and validation of anthropometric-based fat-mass prediction equations using air displacement plethysmography in Mexican infants. *Eur J Clin Nutr.* 2023 Jul;77(7):748-756. doi: 10.1038/s41430-023-01285-9. Epub 2023 Apr 13. PMID: 37055482; PMCID: PMC10335931.
3. LYONS-REID J, et al. Prediction of fat-free mass in a multi-ethnic cohort of infants using bioelectrical impedance: validation against the PEA POD. *Front Nutr.* 2022 OCT;13;9:980790. DOI: 10.3389/FNUT.2022.980790. ecollection 2022. PMID: 36313113 PMCID: PMC9606768
4. GOPALAKRISHNAMOORTHY M, et al. Anthropometric models to estimate fat mass at 3 days, 15 and 54 weeks. *Pediatr Obes.* 2022 Mar 17(3):e12855. doi: 10.1111/ijpo.12855. Epub 2021 Sep 24. PMID: 34558804 PMCID: PMC8821135
5. BANTING S, et al. Estimation of neonatal body fat percentage predicts neonatal hypothermia better than birthweight centile. *J Matern Fetal Neonatal Med* 2022 Feb 1;1-8. doi: 10.1080/14767058.2022.2032634. Online ahead of print. PMID: 35105273
6. OLGA L, et al. Anthropometry-based prediction of body composition in early infancy compared to air-displacement plethysmography. *Pediatr Obes.* 2021 Nov;16(11):e12818. doi: 10.1111/ijpo.12818. Epub 2021 Jun 10. PMID: 34114363

7. NAGEL E, et al. Weight for length measures may not accurately reflect adiposity in preterm infants born appropriate for gestational age during hospitalisation or after discharge from the neonatal intensive care unit. *Pediatr Obes.* 2021 May;16(5):e12744. doi: 10.1111/ijpo.12744. Epub 2020 Nov 3. PMID: 33140910; PMCID: PMC8026714
8. JAYASINGHE S, et al. Anthropometry-based prediction of body fat in infants from birth to 6 months: the Baby-bod study. *Eur J Clin Nutr.* 2021 Apr;75(4):715-723. doi: 10.1038/s41430-020-00768-3. Epub 2020 Oct 14. PMID: 33057177
9. YUMANI DFJ, et al. A comparative study using dual-energy X-ray absorptiometry, air displacement plethysmography, and skinfolds to assess fat mass in preterms at term equivalent age. *Eur J Pediatr.* 2021 Mar;180(3):919-927. doi: 10.1007/s00431-020-03812-3. Epub 2020 Oct 1. PMID: 33006007; PMCID: PMC7886735
10. NAGEL E, et al. Can ultrasound measures of muscle and adipose tissue thickness predict body composition of premature infants in the neonatal intensive care unit? *J Parenter Enteral Nutr.* 2021 Feb;45(2):323-330. doi: 10.1002/jpen.1829. Epub 2020 Apr 7. PMID: 32255211
11. KOUWENHOVEN SMP, et al. Methods to Assess Fat Mass in Infants and Young Children: A Comparative Study Using Skinfold Thickness and Air-Displacement Plethysmography. *Life (Basel).* 2021 Jan 20;11(2):75. doi: 10.3390/life11020075. PMID: 33498272; PMCID: PMC7909249
12. MARANO D, et al. Evaluation of anthropometric equations developed to estimate neonates' body composition: A systematic review. *Cien Saude Colet.* 2020 Jul 8;25(7):2711-2720. doi: 10.1590/1413-81232020257.26982018. PMID: 32667553
13. HAWKES C, et al. The relationship between IGF-I and -II concentrations and body composition at birth and over the first 2 months. *Pediatr Res.* 2019 Apr;85(5):687-692. doi: 10.1038/s41390-019-0331-x. Epub 2019 Feb 11. PMID: 30745570; PMCID: PMC6435390
14. RODRIGUEZ-CANO A, et al. Anthropometric and clinical correlates of fat mass in healthy term infants at 6 months of age. *BMC Pediatr.* 2019 Feb 18;19(1):60. doi: 10.1186/s12887-019-1430-x. PMID: 30777039; PMCID: PMC6378706
15. JOSEFSON J, et al. Fat mass estimation in neonates: Anthropometric models compared with air displacement plethysmography. *Br J Nutr.* 2019 Feb;121(3):285-290. doi: 10.1017/S0007114518003355. Epub 2018 Nov 16. PMID: 30444206; PMCID: PMC7009914
16. ROY S, et al. Body Mass Index is a better indicator of body composition than weight-for-length at age 1 month. *J Pediatr.* 2019 Jan;204:77-83.e1. doi: 10.1016/j.jpeds.2018.08.007. Epub 2018 Sep 26. PMID: 30268397; PMCID: PMC6309630
17. HUVANANDANA J, et al. An anthropometric approach to characterizing neonatal morbidity and body composition, using air displacement plethysmography as a criterion method. *PLoS One.* 2018 Mar 30;13(3):e0195193. doi: 10.1371/journal.pone.0195193. PMID: 29601596; PMCID: PMC5877876
18. CHEN L, et al. Which anthropometric measures best reflect neonatal adiposity? *Int J Obes (Lond).* 2018 Mar;42(3):501-506. doi: 10.1038/ijo.2017.250. Epub 2017 Oct 9. PMID: 28990589; PMCID: PMC5862425
19. LIOTTO N, et al. Can basic characteristics estimate body composition in early infancy? *J Pediatr Gastroenterol Nutr.* 2018 Mar;66(3):e76-e80. doi: 10.1097/MPG.0000000000001758. PMID: 28953532
20. PERNG W, et al. An observational cohort study of weight- and length-derived anthropometric indicators with body composition at birth and 5 mo: The Healthy Start Study. *Am J Clin Nutr.* 2017 Aug;106(2):559-567. doi: 10.3945/ajcn.116.149617. Epub 2017 Jun 28. PMID: 28659296; PMCID: PMC5525117
21. RAMEL S, et al. Do anthropometric measures accurately reflect body composition in preterm infants? *Pediatr Obes.* 2017 Aug;12 Suppl 1:72-77. doi: 10.1111/ijpo.12181. Epub 2016 Sep 16. PMID: 27635625
22. ROELANTS J, et al. Foetal fractional thigh volume: an early 3D ultrasound marker of neonatal adiposity. *Pediatr Obes.* 2017 Aug;12 Suppl 1:65-71. doi: 10.1111/ijpo.12231. Epub 2017 Jul 28. PMID: 28752648
23. CAUBLE J, et al. Validity of anthropometric equations to estimate infant fat mass at birth and in early infancy. *BMC Pediatr.* 2017 Mar 27;17(1):88. doi: 10.1186/s12887-017-0844-6. Erratum in: *BMC Pediatr.* 2020 Feb 28;20(1):92. PMID: 28347278; PMCID: PMC5368988
24. LARCADE J, et al. Estimation of fat-free mass at discharge in preterm infants fed with optimized feeding regimen. *J Pediatr Gastroenterol Nutr.* 2017 Jan;64(1):115-118. doi: 10.1097/MPG.0000000000001261. PMID: 27149252

25. HAWKES C, et al. Body Composition within the first 3 months: Optimized correction for length and correlation with BMI at 2 years. *Horm Res Paediatr*. 2016;86(3):178-187. doi: 10.1159/000448659. Epub 2016 Aug 25. PMID: 27560149
26. WROTTESLEY S, et al. A comparison of body composition estimates using dual-energy X-ray absorptiometry and air-displacement plethysmography in South African neonates. *Eur J Clin Nutr*. 2016 Nov;70(11):1254-1258. doi: 10.1038/ejcn.2016.91. Epub 2016 Jun 1. PMID: 27245207
27. KIGER J, et al. Preterm infant body composition cannot be accurately determined by weight and length. *J Neonatal Perinatal Med*. 2016 Sep 16;9(3):285-90. doi: 10.3233/NPM-16915125. PMID: 27589548
28. MOORE G, et al. Can Fetal Limb Soft Tissue Measurements in the Third Trimester Predict Neonatal Adiposity? *J Ultrasound Med*. 2016 Sep;35(9):1915-24. doi: 10.7863/ultra.15.06028. Epub 2016 Jul 14. PMID: 27417735
29. BARBOUR L, et al. Striking differences in estimates of infant adiposity by new and old DXA software, PEAPOD and skin-folds at 2 weeks and 1 year of life. *Pediatr Obes*. 2016 Aug;11(4):264-71. doi: 10.1111/ijpo.12055. Epub 2015 Jul 22. PMID: 26198967; PMCID: PMC4762753
30. RAUSCH I, et al. Reproducibility of MRI Dixon-based attenuation correction in combined PET/MR with applications for lean body mass estimation. *J Nucl Med*. 2016 Jul;57(7):1096-101. doi: 10.2967/jnumed.115.168294. Epub 2016 Feb 25. PMID: 26917707
31. TINT M, et al. Estimation of fat-free mass in Asian neonates using bioelectrical impedance analysis. *Br J Nutr*. 2016 Mar 28;115(6):1033-42. doi: 10.1017/S0007114515005486. Epub 2016 Feb 9. PMID: 26856420; PMCID: PMC4820031
32. WILBÆK R, et al. Calibration of bioelectrical impedance analysis for body composition assessment in Ethiopian infants using air-displacement plethysmography. *Eur J Clin Nutr*. 2015 Oct;69(10):1099-104. doi: 10.1038/ejcn.2015.51. Epub 2015 Apr 1. PMID: 25828629
33. DALY-WOLF K, et al. Mid-arm circumference is a reliable method to estimate adiposity in preterm and term infants. *Pediatr Res*. 2015 Sep;78(3):336-41. doi: 10.1038/pr.2015.103. Epub 2015 May 28. PMID: 26020147
34. GRIJALVA-ETERNOD C, et al. Midupper arm circumference and weight-for-length z scores have different associations with body composition: Evidence from a cohort of Ethiopian infants. *Am J Clin Nutr*. 2015 Sep;102(3):593-9. doi: 10.3945/ajcn.114.106419. Epub 2015 Jul 29. PMID: 26224296
35. O'CONNOR C, et al. Fetal subcutaneous tissue measurements in pregnancy as a predictor of neonatal total body composition. *Prenat Diagn*. 2014 Oct;34(10):952-5. doi: 10.1002/pd.4400. Epub 2014 Jun 9. PMID: 24788484
36. DE CUNTO A, et al. Can body mass index accurately predict adiposity in newborns? *Arch Dis Child Fetal Neonatal*. 2014 May;99(3):F238-9. doi: 10.1136/archdischild-2013-305386. Epub 2013 Dec 3. PMID: 24302686
37. ARIS I, et al. Body fat in Singaporean infants: Development of body fat prediction equations in Asian newborns. *Eur J Clin Nutr*. 2013 Sep;67(9):922-7. doi: 10.1038/ejcn.2013.69. Epub 2013 Apr 3. PMID: 23549200
38. LINGWOOD B, et al. Prediction of fat-free mass and percentage of body fat in neonates using bioelectrical impedance analysis and anthropometric measures: Validation against the PEA POD. *Br J Nutr*. 2012 May;107(10):1545-52. doi: 10.1017/S0007114511004624. Epub 2011 Sep 15. PMID: 21917194
39. DEIERLEIN A, et al. An anthropometric model to estimate neonatal fat mass using air displacement plethysmography. *Nutr Metab (Lond)*. 2012 Mar 21;9:21. doi: 10.1186/1743-7075-9-21. PMID: 22436534; PMCID: PMC3348023.
40. LEE W, et al. Fetal growth parameters and birth weight: their relationship to neonatal body composition. *Ultrasound Obstet Gynecol*. 2009 Apr;33(4):441-6. doi: 10.1002/uog.6317. PMID: 19253324; PMCID: PMC3549427

## Review

1. COUTO E, et al. Predictive models of newborn body composition: a systematic review. *Rev Paul Pediatr*. 2023 Mar 13;41:e2020365. doi: 10.1590/1984-0462/2023/41/2020365. PMID: 36921160; PMCID: PMC10014017.

2. YUMANI D, et al. Body composition in preterm infants: a systematic review on measurement methods. *Pediatr Res*. 2023 Apr;93(5):1120-1140. doi: 10.1038/s41390-022-02262-x. Epub 2022 Aug 22. PMID: 35995939.
3. BELL K, et al. Body composition measurement for the preterm neonate: using a clinical utility framework to translate research tools into clinical care. *J Perinatol*. 2022 Nov;42(11):1550-1555. doi: 10.1038/s41372-022-01529-9. Epub 2022 Oct 6. PMID: 36203085; PMCID: PMC9617782.
4. WHYTE K, et al. 1.2.3 Technical Measurements of Body Composition Assessment. *World Rev Nutr Diet*. 2022;124:23-30. doi: 10.1159/000517192. Epub 2022 Mar 3. PMID: 35240646.
5. RALLIS D, et al. A systematic review indicates an association between birth weight and body fat in childhood. *Acta Paediatr*. 2021 Jul;110(7):2023-2039. doi: 10.1111/apa.15834. Epub 2021 Mar 15. PMID: 33682216
6. GALLAGHER D, et al. Body composition measurements from birth through 5 years: Challenges, gaps, and existing & emerging technologies - A National Institutes of Health workshop. *Obes Rev*. 2020 Aug;21(8):e13033. doi: 10.1111/obr.13033. Epub 2020 Apr 20. PMID: 32314544
7. CERASANI J, et al. Human milk feeding and preterm infants' growth and body composition: A literature review. *Nutrients*. 2020 Apr 21;12(4):1155. doi: 10.3390/nu12041155. PMID: 32326178; PMCID: PMC7230190
8. ANDREWS E, et al. Measuring body composition in the preterm infant: Evidence base and practicalities. *Clin Nutr*. 2019 Dec;38(6):2521-2530. doi: 10.1016/j.clnu.2018.12.033. Epub 2019 Jan 12. PMID: 30737045
9. BELFORT M, et al. NICU Diet, physical growth and nutrient accretion, and preterm infant brain development. *Neoreviews*. 2019 Jul;20(7):e385-e396. doi: 10.1542/neo.20-7-e385. PMID: 31261105
10. STRYDOM K, et al. Factors affecting body composition in preterm infants: Assessment techniques and nutritional interventions. *Pediatr Neonatol*. 2019 Apr;60(2):121-128. doi: 10.1016/j.pedneo.2017.10.007. Epub 2017 Oct 28. PMID: 29239827
11. MAZAHERY H, et al. Air displacement plethysmography (Pea Pod) in full-term and pre-term infants: a comprehensive review of accuracy, reproducibility, and practical challenges. *Matern Health Neonatol Perinatol*. 2018 Jun 20;4:12. doi: 10.1186/s40748-018-0079-z. PMID: 29951209; PMCID: PMC6011189
12. ROELANTS J, et al. Prenatal markers of neonatal fat mass: A systematic review. *Clin Nutr*. 2016 Oct;35(5):995-1007. doi: 10.1016/j.clnu.2015.09.003. Epub 2015 Sep 25. PMID: 26499032
13. TORO-RAMOS T, et al. Body composition during fetal development and infancy through the age of 5 years. *Eur J Clin Nutr*. 2015 Dec;69(12):1279-89. doi: 10.1038/ejcn.2015.117. Epub 2015 Aug 5. PMID: 26242725; PMCID: PMC4680980
14. RICE M, et al. Neonatal body composition: Measuring lean mass as a tool to guide nutrition management in the neonate. *Nutr Clin Pract*. 2015 Oct;30(5):625-32. doi: 10.1177/0884533615578917. Epub 2015 Apr 23. PMID: 25908606
15. ZANINI RDE V, et al. Body fat in children measured by DXA, air-displacement plethysmography, TBW and multicomponent models: A systematic review. *Matern Child Health J*. 2015 Jul;19(7):1567-73. doi: 10.1007/s10995-015-1666-5. PMID: 25636644
16. FIELDS D, et al. Air displacement plethysmography: Cradle to grave. *Nutr Clin Pract*. 2015 Apr;30(2):219-26. doi: 10.1177/0884533615572443. Epub 2015 Mar 11. PMID: 25761768
17. HORAN M, et al. Methodologies to assess paediatric adiposity. *J Med Sci*. 2015 Mar;184(1):53-68. doi: 10.1007/s11845-014-1124-1. Epub 2014 May 4. PMID: 24791970
18. DEMERATH W, et al. Body composition assessment in the infant. *Am J Hum Biol*. 2014 May-Jun;26(3):291-304. doi: 10.1002/ajhb.22500. Epub 2014 Jan 15. PMID: 24424686; PMCID: PMC5761669
19. WELLS J, et al. Toward body composition reference data for infants, children, and adolescents. *Adv Nutr*. 2014 May 14;5(3):320S-9S. doi: 10.3945/an.113.005371. PMID: 24829484; PMCID: PMC4013189
20. CORPELEIJN W, et al. Optimal growth of preterm infants. *World Rev Nutr Diet*. 2013;106:149-55. doi: 10.1159/000342584. Epub 2013 Feb 11. PMID: 23428694

21. WARD L, et al. Assessing early growth and adiposity: Report from an EarlyNutrition Academy workshop. *Ann Nutr Metab.* 2013;63(1-2):120-30. doi: 10.1159/000350702. Epub 2013 Aug 20. PMID: 23969405
22. LI C, et al. Infant body composition in the PEA POD® era: What have we learned and where do we go from here? *J Dev Orig Health Dis.* 2013 Apr;4(2):116-20. doi: 10.1017/S2040174412000657. PMID: 25054677
23. WELLS J. Body composition in infants: Evidence for developmental programming and techniques for measurement. *Rev Endocr Metab Disord.* 2012 Jun;13(2):93-101. doi: 10.1007/s11154-012-9213-9. PMID: 22418619 13:93-101 (2012)
24. BARACOS V, et al. Advances in the science and application of body composition measurement. *JPEN J Parenter Enteral Nutr.* 2012 Jan;36(1):96-107. doi: 10.1177/0148607111417448. PMID: 22235108; PMCID: PMC4422066
25. ELLIS K. Body composition in infancy: Impact on health later in life. *Nestlé Nutr Inst Workshop Ser Pediatr Program.* 2010;65:213-20; discussion 221-4. doi: 10.1159/000281168. Epub 2010 Feb 1. PMID: 20139684.
26. RUBIN L. Postnatal growth in preterm infants: Too small, too big, or just right? *J Pediatr.* 2009 Apr;154(4):473-5. doi: 10.1016/j.jpeds.2008.12.041. PMID: 19324213
27. ROGGERO P, et al. Measuring the body composition of preterm and term neonates: From research to clinical applications. *J Pediatr Gastroenterol Nutr.* 2007 Dec;45 Suppl 3:S159-62. doi: 10.1097/01.mpg.0000302964.85922.1a. Erratum in: *J Pediatr Gastroenterol Nutr.* 2009 Jan;48(1):121-2. PMID: 18185084
28. WELLS J, et al. Programming of body composition by early growth and nutrition. *Pro Nutr Soc.* 2007 Aug;66(3):423-34. doi: 10.1017/S0029665107005691. PMID: 17637095 66:423-434 (2007)
29. ELLIS K. Evaluation of body composition in neonates and infants. *Semin Fetal Neonatal Med.* 2007 Feb;12(1):87-91. doi: 10.1016/j.siny.2006.10.011. Epub 2006 Dec 14. PMID: 17174616
30. RIGO J. Body composition during the first year of life. *Nestlé Nutr Inst Workshop Ser Pediatr Program.* 2006;58:65-76; discussion 76-8. doi: 10.1159/000095021. PMID: 16902326, 58:65-78 (2006)

## Normative Body Composition

1. MURPHY-ALFORD, A, et al. Body composition reference charts for infants from birth to 24 months: Multicenter Infant Body Composition Reference Study. *Am J Clin Nutr.* 2023 Jun;117(6):1262-1269. doi: 10.1016/j.ajcnut.2023.02.012. Epub 2023 May 11. PMID: 37270290.
2. WONG R, et al. Infant Body Composition in an Asian Pacific Islander Population. *J Racial Ethn Health Disparities.* 2022 Nov 10. doi: 10.1007/s40615-022-01444-x. Epub ahead of print. PMID: 36357640.
3. YOUSUF E, et al. Growth and body composition trajectories in infants meeting the WHO growth standards study requirements. *Pediatr Res.* 2022 Mar 16. doi: 10.1038/s41390-022-02005-y. Online ahead of print. PMID: 35296784
4. JAYASINGHE S, et al. WHO Child Growth Standards in context: The Baby-bod Project - Observational study in Tasmania. *BMJ Paediatr Open.* 2021 Jun 16;5(1):e001123. doi: 10.1136/bmjpo-2021-001123. PMID: 34222680; PMCID: PMC8211047
5. DE FLUITER K, et al. Longitudinal body composition assessment in healthy term-born infants until 2 years of age using ADP and DXA with vacuum cushion. *Eur J Clin Nutr.* 2021 May;40(5):3401-3408. doi: 10.1016/j.clnu.2020.11.024. Epub 2020 Nov 24. PMID: 33309161
6. LUTZ T, et al. A hospital-based cohort study of gender and gestational age-specific body fat percentage at birth. *Pediatr Res.* 2021 Jan;89(1):231-237. doi: 10.1038/s41390-020-0879-5. Epub 2020 Apr 14. PMID: 32289811
7. KURIYAN R, et al. The thin but fat phenotype is uncommon at birth in Indian babies. *J Nutr.* 2020 Apr 1;150(4):826-832. doi: 10.1093/jn/nxz305. PMID: 31858112
8. WIECHERS C, et al. Neonatal body composition by air displacement plethysmography in healthy term singletons: A systematic review. *BMC Pediatr.* 2019 Dec 12;19(1):489. doi: 10.1186/s12887-019-1867-y. PMID: 31830946; PMCID: PMC6907141
9. WIECHERS C, et al. Neonatal body composition: Cross-sectional study in healthy term singletons in Germany. *BMC Pediatr.* 2019 Dec 12;19(1):488. doi: 10.1186/s12887-019-1837-4. PMID: 31830928; PMCID: PMC6907196.

10. ALEXANDER T, et al. Body composition of New Zealand-born term babies differs by ethnicity, gestational age and sex. *Early Hum Dev.* 2019 Nov 15;140:104924. doi: 10.1016/j.earlhumdev.2019.104924. Epub ahead of print. PMID: 31739266
11. FORSUM E, et al. Fat and fat-free mass of healthy Swedish children show tracking during early life, but there are differences. *Acta Paediatr.* 2019 Sep;108(9):1704-1708. doi: 10.1111/apa.14771. Epub 2019 Mar 25. PMID: 30830968
12. NORRIS T, et al. New charts for the assessment of body composition, according to air-displacement plethysmography, at birth and across the first 6 mo of life. *Am J Clin Nutr.* 2019 May 1;109(5):1353-1360. doi: 10.1093/ajcn/nqy377. PMID: 31051509
13. HENRICKSSON P, et al. Longitudinal assessment of body composition in healthy Swedish children from 1 week until 4 years of age. *Eur J Clin Nutr.* 2017 Aug 23. doi: 10.1038/ejcn.2017.125. Epub ahead of print. PMID: 28832576
14. VILLAR J, et al. Body composition at birth and its relationship with neonatal anthropometric ratios: The newborn body composition study of the INTERGROWTH-21st project. *Pediatr Res.* 2017 Aug;82(2):305-316. doi: 10.1038/pr.2017.52. Epub 2017 May 31. PMID: 28445454; PMCID: PMC5605677
15. SCHEURER J, et al. Body composition trajectories from infancy to preschool in children born premature versus full-term. *J Pediatr Gastroenterol Nutr.* 2017 Jun;64(6):e147-e153. doi: 10.1097/MPG.0000000000001494. PMID: 28045768
16. DEMERATH E, et al. New body composition reference charts for preterm infants. *Am J Clin Nutr.* 2017 Jan;105(1):70-77. doi: 10.3945/ajcn.116.138248. Epub 2016 Nov 2. PMID: 27806978
17. RAMEL S, et al. Body composition at birth in preterm infants between 30 and 36 weeks gestation. *Pediatr Obes.* 2015 Feb;10(1):45-51. doi: 10.1111/j.2047-6310.2013.00215.x. Epub 2014 Jan 27. PMID: 24470220
18. PAVIOTTI G, et al. Longitudinal growth and body composition of twins versus singleton in the first month of life. *Scientific World Journal.* 2013 Dec 24;2013:108189. doi: 10.1155/2013/108189. PMID: 24453786; PMCID: PMC3886583
19. CARBERRY A, et al. Is body fat percentage a better measure of undernutrition in newborns than birth weight percentiles? *Pediatr Res.* 2013 Dec;74(6):730-6. doi: 10.1038/pr.2013.156. Epub 2013 Sep 3. PMID: 24002331
20. ANDERSEN G, et al. Body composition from birth to 6 mo. of age in Ethiopian infants: Reference data obtained by air-displacement plethysmography. *Am J Clin Nutr.* 2013 Oct;98(4):885-94. doi: 10.3945/ajcn.113.063032. Epub 2013 Aug 28. PMID: 23985805
21. JOHNSON M, et al. Preterm birth and body composition at term equivalent age: A systematic review and meta-analysis. *Pediatrics.* 2012 Sep;130(3):e640-9. doi: 10.1542/peds.2011-3379. Epub 2012 Aug 13. PMID: 22891222
22. LAW T, et al. Customized versus population-based growth curves: Prediction of low body fat percent at term corrected gestational age following preterm birth. *J Matern Fetal Neonatal Med.* 2012 Jul;25(7):1142-7. doi: 10.3109/14767058.2011.625459. Epub 2012 Apr 2. PMID: 21939292
23. FIELDS D, et al. Longitudinal body composition data in exclusively breast-fed infants: A multicenter study. *Obesity (Silver Spring).* 2011 Sep;19(9):1887-91. doi: 10.1038/oby.2011.11. Epub 2011 Feb 10. PMID: 21311509 19(9):1887-1891 (2011)
24. HAWKES C, et al. Gender-and gestational age-specific body fat percentage at birth. *Pediatrics.* 2011 Sep;128(3):e645-51. doi: 10.1542/peds.2010-3856. Epub 2011 Aug 8. PMID: 21824882
25. ROGGERO P, et al. Quality of growth in exclusively breast-fed infants in the first six months of life: An Italian study. *Pediatr Res.* 2010 Dec;68(6):542-4. doi: 10.1203/PDR.0b013e3181f85a20. PMID: 20736880
26. CARBERRY A, et al. Body composition from birth to 4.5 months in infants born to non-obese women. *Pediatr Res* 2010 Jul;68(1):84-8. doi: 10.1203/PDR.0b013e3181df5421. PMID: 20351656

### Body Composition Observation in Preterm Infants

1. THAJER A, et al. The Impacts of Single Preterm Human Donor Milk Compared to Mother's Own Milk on Growth and Body Composition. *Nutrients.* 2023;15(7):1578. Published 2023 Mar 24. doi:10.3390/nu15071578

2. MORRIS E, et al. Preterm infant body composition, working memory, and temperament. *Infant Behav Dev.* 2023 Feb;70:101808. doi: 10.1016/j.infbeh.2022.101808. Epub 2023 Jan 5. PMID: 36610269.
3. BELL K, et al. Associations of body composition with regional brain volumes and white matter microstructure in very preterm infants. *Arch Dis Child Fetal Neonatal Ed.* 2022 Sep;107(5):533-538. doi: 10.1136/archdischild-2021-321653. Epub 2022 Jan 20. PMID: 35058276 PMCID: PMC929669
4. LIMA P, et al. Energy expenditure and body composition in infants with bronchopulmonary dysplasia at term age. *European Journal of Pediatrics.* 2022 Aug;181(8):3039-3047. doi: 10.1007/s00431-022-04517-5. Epub 2022 Jun 4. PMID: 35661246
5. LACH L, et al. Body Composition and "Catch-Up" Fat Growth in Healthy Small for Gestational Age Preterm Infants and Neurodevelopmental Outcomes. *Nutrients.* 2022 Jul 25;14(15):3051. doi: 10.3390/nu14153051. PMID: 35893903 PMCID: PMC9332383
6. MCNELIS K. et al. Body Composition of Infants With Congenital Gastroschisis. *J Pediatr Gastroenterol Nutr.* 2022 Feb 1;74(2):e35-e38. doi: 10.1097/MPG.0000000000003288. PMID: 34406999
7. NAGEL E, et al. Ultrasound measurements of abdominal muscle thickness are associated with postmenstrual age at full oral feedings in preterm infants: A preliminary study. *Nutr Clin Pract.* 2021 Dec;36(6):1207-1214. doi: 10.1002/ncp.10670. Epub 2021 Jun 8. PMID: 34101241 PMCID: PMC8651832
8. BINDER C, et al. Association between Fat-Free Mass and Brain Size in Extremely Preterm Infants. *Nutrients.* 2021 Nov 24;13(12):4205. doi: 10.3390/nu13124205. PMID: 34959757 PMCID: PMC8708955
9. PERRONE M, et al. Energy Expenditure, Protein Oxidation and Body Composition in a Cohort of Very Low Birth Weight Infants. *Nutrients.* 2021 Nov 6;13(11):3962. doi: 10.3390/nu13113962. PMID: 34836218 PMCID: PMC8620881
10. BUA J, et al. Association between body composition at term equivalent age and Bayley scores at 2 years in preterm infants. *J Perinatol.* 2021 Aug;41(8):1852-1858. doi: 10.1038/s41372-021-01074-x. Epub 2021 May 13. PMID: 33986476
11. YUMANI DFJ, et al. IGF-I, Growth, and Body Composition in Preterm Infants up to Term Equivalent Age. *J Endocr Soc.* 2021 Jun 18;5(7):bvab089. doi: 10.1210/jendso/bvab089. PMID: 34159288; PMCID: PMC8212689
12. BEUNDERS V, et al. Early weight gain trajectories and body composition in infancy in infants born very preterm. *Pediatr Obes.* 2021 Jun;16(6):e12752. doi: 10.1111/ijpo.12752. Epub 2020 Nov 17. PMID: 33205622; PMCID: PMC8244114
13. PAVIOTTI G, et al. Change in body composition of premature infants from parenteral nutrition discontinuation to term equivalent age. *Early Hum Dev.* 2020 Dec;151:105163. doi: 10.1016/j.earlhumdev.2020.105163. Epub 2020 Aug 20. PMID: 32905898
14. SALAS A, et al. Serial assessment of fat and fat-free mass accretion in very preterm infants: A randomized trial. *Pediatr Res.* 2020 Nov;88(5):733-738. doi: 10.1038/s41390-020-1052-x. Epub 2020 Jul 7. PMID: 32634820; PMCID: PMC7581604
15. BRUCKNER M, et al. Extremely preterm infants have a higher fat mass percentage in comparison to very preterm infants at term-equivalent age. *Front Pediatr.* 2020 Mar 10;8:61. doi: 10.3389/fped.2020.00061. PMID: 32219084; PMCID: PMC7078645
16. FORSUM E, et al. Premature birth was not associated with increased body fatness in four-year-old boys and girls. *Acta Paediatr.* 2020 Feb;109(2):327-331. doi: 10.1111/apa.14990. Epub 2019 Sep 16. PMID: 31461786
17. HAMATSCHEK C, et al. Fat and fat-free mass of preterm and term infants from birth to six months: A review of current evidence. *Nutrients.* 2020 Jan 21;12(2):288. doi: 10.3390/nu12020288. PMID: 31973218; PMCID: PMC7070317
18. RAMEL S, et al. Nutrition, illness and body composition in very low birth weight preterm infants: Implications for nutrition and neurocognitive outcomes. *Nutrients.* 2020 Jan 4;12(1):145. doi: 10.3390/nu12010145. PMID: 31947964; PMCID: PMC7019791
19. CHMIELEWSKA A, et al. Lean tissue deficit in preterm infants persists up to 4 months of age: Results from a Swedish longitudinal study. *Neonatology.* 2020;117(1):80-87. doi: 10.1159/000503292. Epub 2019 Dec 10. PMID: 31822002
20. INGOLFSLAND E, et al. Late growth and changes in body composition influence odds of developing retinopathy of prematurity among preterm infants. *Nutrients.* 2019 Dec 27;12(1):78. doi: 10.3390/nu12010078. PMID: 31892145; PMCID: PMC7019301
21. BELL K, et al. Associations of growth and body composition brain size in preterm infants. *J Pediatr.* 2019 Nov;214:20-26.e2. doi: 10.1016/j.jpeds.2019.06.062. Epub 2019 Jul 31. PMID: 31377040

22. MORNIROLI D, et al. Is the body composition development in premature infants associated with a distinctive nuclear magnetic resonance metabolomic profiling of urine? *J Matern Fetal Neonatal Med.* 2019 Jul;32(14):2310-2318. doi: 10.1080/14767058.2018.1432040. Epub 2018 Feb 15. PMID: 29357769
23. MC CLUNAN K, et al. Effect of nutritional intake on the body composition of HIV-exposed and HIV-unexposed preterm and low birth weight infants. *Breastfeed Med.* 2019 Apr;14(3):144-153. doi: 10.1089/bfm.2018.0113. Epub 2019 Feb 4. PMID: 30715911
24. JEROME M, et al. Racial disparities in body composition of preterm infants (P21-049-19). *Curr Dev Nutr.* 2019;3(Suppl 1):nzz041.P21-049-19
25. PIEMONTESE P, et al. The Effect of Human Milk on Modulating the Quality of Growth in Preterm Infants. *Front Pediatr.* 2018 Oct 9;6:291. doi: 10.3389/fped.2018.00291. PMID: 30356677; PMCID: PMC6189304
26. DA SILVA MARTINS A, et al. Growth and body composition in preterm newborns with bronchopulmonary dysplasia: A cohort study. *J Perinat Med.* 2018 Oct 25;46(8):913-918. doi: 10.1515/jpm-2018-0005. PMID: 29949515
27. VILLELA L, et al. Body composition in preterm infants with intrauterine growth restriction: A cohort study. *J Perinat Med.* 2018 Sep 25;46(7):804-810. doi: 10.1515/jpm-2017-0175. PMID: 29408800
28. SCHEURER J, et al. Body Composition Changes from Infancy to 4 Years and Associations with Early Childhood Cognition in Preterm and Full-Term Children. *Neonatology.* 2018;114(2):169-176. doi: 10.1159/000487915. Epub 2018 Jun 13. PMID: 29898453; PMCID: PMC6083858
29. VILLELA L, et al. Growth and body composition of preterm infants less than or equal to 32 weeks: Cohort study. *Early Hum Dev.* 2018 Feb;117:90-95. doi: 10.1016/j.earlhumdev.2018.01.007. Epub 2018 Jan 23. PMID: 29360048
30. ALGOTAR A, et al. Unique patterns of body composition and anthropometric measurements during maturation in neonatal intensive care unit neonates: Opportunities for modifying nutritional therapy and influencing clinical outcomes. *JPEN J Parenter Enteral Nutr.* 2018 Jan;42(1):231-238. doi: 10.1002/jpen.1012. Epub 2017 Dec 13. PMID: 29505146
31. ABERA M, et al. Relation between body composition at birth and child development at 2 years of age: A prospective cohort study among Ethiopian children. *Eur J Clin Nutr.* 2017 Dec;71(12):1411-1417. doi: 10.1038/ejcn.2017.129. Epub 2017 Sep 27. PMID: 28952606
32. PAVIOTTI G, et al. Higher growth, fat and fat-free masses correlate with larger cerebellar volumes in preterm infants at term. *Acta Paediatr.* 2017 Jun;106(6):918-925. doi: 10.1111/apa.13829. Epub 2017 Apr 20. PMID: 28295577
33. PIYASENA C, et al. Dynamic changes in DNA methylation occur during the first year of life in preterm infants. *Front Endocrinol (Lausanne).* 2016 Dec 15;7:158. doi: 10.3389/fendo.2016.00158. Erratum in: *Front Endocrinol (Lausanne).* 2018 Feb 27;9:47. PMID: 28018293; PMCID: PMC5156662
34. RAMEL S, et al. Greater early gains in fat-free mass, but not fat mass, are associated with improved neurodevelopment at 1 year corrected age for prematurity in very low birth weight preterm infants. *J Pediatr.* 2016 Jun;173:108-15. doi: 10.1016/j.jpeds.2016.03.003. Epub 2016 Apr 4. PMID: 27056450
35. GIANNI M, et al. Body composition in late preterm infants according to percentile at birth. *Pediatr Res.* 2016 May;79(5):710-5. doi: 10.1038/pr.2015.273. Epub 2015 Dec 30. PMID: 26717003
36. SCHEURER J, et al. Diminished growth and lower adiposity in hyperglycemic very low birth weight neonates at 4 months corrected age. *J Perinatol.* 2016 Feb;36(2):145-50. doi: 10.1038/jp.2015.154. Epub 2015 Nov 5. PMID: 26540246
37. DE CUNTO A, et al. Impact of surgery for neonatal gastrointestinal diseases on weight and fat mass. *J Pediatr.* 2015 Sep;167(3):568-71. doi: 10.1016/j.jpeds.2015.06.013. Epub 2015 Jul 3. PMID: 26148657
38. ROGGERO P, et al. No relative increase in intra-abdominal adipose tissue in healthy unstressed preterm infants at term. *Neonatology.* 2015;107(1):14-9. doi: 10.1159/000364855. Epub 2014 Oct 4. PMID: 25301293
39. PIYASENA C, et al. Dynamics of DNA methylation at IGF2 in preterm and term infants during the first year of life: An observational study. *Lancet.* 2015 Feb 26;385 Suppl 1:S81. doi: 10.1016/S0140-6736(15)60396-8. PMID: 26312903

40. OLHAGER E, et al. Body composition in late preterm infants in the first 10 days of life and at full term. *Acta Paediatr.* 2014 Jul;103(7):737-43. doi: 10.1111/apa.12632. Epub 2014 May 7. PMID: 24628453
41. SIMON L, et al. Determinants of body composition in preterm infants at the time of hospital discharge. *Am J Clin Nutr.* 2014 Jul;100(1):98-104. doi: 10.3945/ajcn.113.080945. Epub 2014 May 7. PMID: 24808483
42. PFISTER K, et al. Exploratory study of the relationship of fat-free mass to speed of brain processing in preterm infants. *Pediatr Res.* 2013 Nov;74(5):576-83. doi: 10.1038/pr.2013.138. Epub 2013 Aug 13. PMID: 23942556
43. MEYERS J, et al. Potential influence of total parenteral nutrition on body composition at discharge in preterm infants. *J Matern Fetal Neonatal Med.* 2013 Oct;26(15):1548-53. doi: 10.3109/14767058.2013.793663. Epub 2013 May 7. PMID: 23578184
44. LIOTTO N, et al. Growth and body composition changes in late preterm infants in the first months of life. *Pediatr Med Chir.* 2013 Jul-Aug;35(4):172-6. Italian. doi: 10.4081/pmc.2013.37. PMID: 24245099
45. PIEMONTESE P, et al. Effect of prematurity on fat mass distribution and blood pressure at prepubertal age: A follow-up study. *Pediatr Med Chir.* 2013 Jul-Aug;35(4):166-71. Italian. doi: 10.4081/pmc.2013.36. PMID: 24245098
46. SIMON L, et al. Effect of sex and gestational age on neonatal body composition. *Br J Nutr.* 2013 Mar 28;109(6):1105-8. doi: 10.1017/S0007114512002991. Epub 2012 Jul 12. PMID: 22784704
47. BARANOV A, et al. Evaluation of premature infants' nutritional status by air plethysmography: First Russian prospective study. *Vestn Ross Akad Med Nauk.* 2013;(4):10-6. Russian. doi: 10.15690/vramn.v68i4.605. PMID: 24003716
48. GIANNI L, et al. Postnatal catch-up fat after late preterm birth. *Pediatr Res.* 2012 Dec;72(6):637-40. doi: 10.1038/pr.2012.128. Epub 2012 Sep 25. PMID: 23011446
49. TARONI F, et al. Intra-abdominal adiposity in preterm infants: An explorative study. *Pediatr Med Chir.* 2012 Nov-Dec;34(6):283-6. Italian. doi: 10.4081/pmc.2012.52. PMID: 24364134
50. LAW T, et al. Ultrasound assessment of intrauterine growth restriction: Relationship to neonatal body composition. *Am J Obstet Gynecol.* 2011 Sep;205(3):255.e1-6. doi: 10.1016/j.ajog.2011.06.027. Epub 2011 Jun 15. PMID: 22071054
51. RAMEL S, et al. Body composition changes in preterm infants following hospital discharge: Comparison with term infants. *J Pediatr Gastroenterol Nutr.* 2011 Sep;53(3):333-8. doi: 10.1097/MPG.0b013e3182243aa7. PMID: 21602717; PMCID: PMC7680641
52. GIANNI M, et al. Adiposity in small for gestational age preterm infants assessed at term equivalent age. *Arch Dis Child Fetal Neonatal Ed.* 2009 Sep;94(5):F368-72. doi: 10.1136/adc.2008.153163. Epub 2009 May 19. PMID: 19457875
53. ROGGERO P, et al. Is term newborn body composition being achieved postnatally in preterm infants? *Early Hum Dev.* 2009 Jun;85(6):349-52. doi: 10.1016/j.earlhumdev.2008.12.011. Epub 2009 Jan 21. PMID: 19162413
54. TARONI F, et al. Body composition in small for gestational age newborns. *Pediatr Med Chir.* 2008 Nov-Dec;30(6):296-301. Italian. PMID: 19431952
55. ROGGERO P, et al. Postnatal growth failure in preterm infants: Recovery of growth and body composition after term. *Early Hum Dev.* 2008 Aug;84(8):555-9. doi: 10.1016/j.earlhumdev.2008.01.012. Epub 2008 Mar 10. PMID: 18329828

### Body Composition Observation in Term Infants

1. MARTINS B, et al. Early growth and body composition assessed by air displacement plethysmography in infants born with simple gastroschisis. *J Pediatr Surg.* 2023;58(4):741-746. doi:10.1016/j.jpedsurg.2022.11.005
2. COHEN C, et al. Body composition trajectories from birth to 5 years and hepatic fat in early childhood. *Am J Clin Nutr.* 2022 Oct 6;116(4):1010-1018. doi: 10.1093/ajcn/nqac168. PMID: 36055960 PMCID: PMC9535524
3. ONG Y, et al. Newborn body composition and child cardiovascular risk markers: a prospective multi-ethnic Asian cohort study. *Int J Epidemiol.* 2022 Jul 30;dyac154. doi: 10.1093/ije/dyac154. Online ahead of print. PMID: 35906917

4. SANDBORG J, et al. The effects of a lifestyle intervention (the HealthyMoms app) during pregnancy on infant body composition: Secondary outcome analysis from a randomized controlled trial. *Pediatr Obes*. 2022 Jun;17(6):e12894. doi: 10.1111/ijpo.12894. PMID: 35106942 PMCID: PMC9285397
5. TAN K, et al. The Kynurenine Pathway Metabolites in Cord Blood Positively Correlate With Early Childhood Adiposity. *J Clin Endocrinol Metab*. 2022 May 17;107(6):e2464-e2473. doi: 10.1210/clinem/dgac078. PMID: 35150259 PMCID: PMC9113811
6. HESSE H, et al. Differences in body composition and growth persist postnatally in fetuses diagnosed with severe compared to mild fetal growth restriction. *J Neonatal Perinatal Med*. 2022;15(3):589-598. doi: 10.3233/NPM-210872. PMID: 35342050
7. CORR TE, et al. Body composition during the first 4 months in infants affected by neonatal abstinence syndrome: a pilot study. *J Dev Orig Health Dis*. 2022 Feb;13(1):120-127. doi: 10.1017/S2040174421000052. Epub 2021 Mar 2. PMID: 33650484
8. VAN BEIJSTERVELDT IALP, Fat mass and fat-free mass track from infancy to childhood: New insights in body composition programming in early life. *Obesity (Silver Spring)*. 2021 Nov;29(11):1899-1906. doi: 10.1002/oby.23271. Epub 2021 Sep 21. PMID: 34549538
9. SAUDER KA, et al. Fat Mass Accretion from Birth to 5 Years and Metabolic Homeostasis in Childhood: the Healthy Start Study. *J Clin Endocrinol Metab*. 2021 May 13;106(6):1684-1691. doi: 10.1210/clinem/dgab115. PMID: 33616653; PMCID: PMC8118576
10. MOORE BF, et al. Joint effects of ambient air pollution and maternal smoking on neonatal adiposity and childhood BMI trajectories in the Healthy Start study. *Environ Epidemiol*. 2021 May 5;5(3):e142. doi: 10.1097/EE9.000000000000142. PMID: 34131612; PMCID: PMC8196098
11. DE FLUITER K, et al. Longitudinal human milk macronutrients, body composition and infant appetite during early life. *Clin Nutr*. 2021 May;40(5):3401-3408. doi: 10.1016/j.clnu.2020.11.024. Epub 2020 Nov 24. PMID: 33309161
12. DE FLUITER K, et al. Association between fat mass in early life and later fat mass trajectories. *JAMA Pediatr*. 2020 Dec 1;174(12):1141-1148. doi: 10.1001/jamapediatrics.2020.2673. PMID: 32804197; PMCID: PMC7432277
13. HENRIKSSON P, et al. DNA methylation in infants with low and high body fatness. *BMC Genomics*. 2020 Nov 9;21(1):769. doi: 10.1186/s12864-020-07169-7. PMID: 33167873; PMCID: PMC7654595
14. PEREIRA-DA-SILVA L, et al. Evolution of resting energy expenditure, respiratory quotient, and adiposity in infants recovering from corrective surgery of major congenital gastrointestinal tract anomalies: A cohort study. *Nutrients*. 2020 Oct 11;12(10):3093. doi: 10.3390/nu12103093. PMID: 33050623; PMCID: PMC7599456
15. RODRÍGUEZ-CANO A, et al. Higher fat mass and fat mass accretion during the first six months of life in exclusively breastfed infants. *Pediatr Res*. 2020 Feb;87(3):588-594. doi: 10.1038/s41390-019-0542-1. Epub 2019 Aug 21. PMID: 31434104; PMCID: PMC7033039
16. KADAKIA R, et al. Association of cord blood methylation with neonatal leptin: An epigenome wide association study. *PLoS One*. 2019 Dec 18;14(12):e0226555. doi: 10.1371/journal.pone.0226555. PMID: 31851703; PMCID: PMC6919608
17. SMITH H, et al. Difference between body composition of formula- and breastfed infants at birth. *J Dev Orig Health Dis*. 2019 Dec;10(6):616-620. doi: 10.1017/S2040174419000187. Epub 2019 May 28. PMID: 31134881
18. DAVIS S, et al. Sex differences in infant body composition emerge in the first 5 months of life. *J Pediatr Endocrinol Metab*. 2019 Nov 26;32(11):1235-1239. doi: 10.1515/jpem-2019-0243. PMID: 31483758; PMCID: PMC6851433
19. WIBAEK R, et al. Associations of fat mass and fat-free mass accretion in infancy with body composition and cardometabolic risk markers at 5 years: The Ethiopian iABC birth cohort study. *PLoS Med*. 2019 Aug 20;16(8):e1002888. doi: 10.1371/journal.pmed.1002888. PMID: 31430287; PMCID: PMC6701744
20. LARSSON A, et al. Body composition and growth in full-term small for gestational age and large for gestational age Swedish infants assessed with air displacement plethysmography at birth and at 3-4 months of age. *PLoS One*. 2019 May 15;14(5):e0207978. doi: 10.1371/journal.pone.0207978. PMID: 31091240; PMCID: PMC6519902
21. EUCLYDES V, et al. Cord blood concentrations of leptin, zinc- $\alpha$ 2-glycoprotein, and adiponectin, and adiposity gain during the first 3 mo of life. *Nutrition*. 2018 Oct;54:89-93. doi: 10.1016/j.nut.2018.02.012. Epub 2018 Mar 22. PMID: 29758496

22. ADMASSU B, et al. Body composition during early infancy and its relation with body composition at 4 years of age in Jimma, an Ethiopian prospective cohort study. *Nutr Diabetes*. 2018 Sep 7;8(1):46. doi: 10.1038/s41387-018-0056-7. PMID: 30190452; PMCID: PMC6127223
23. KADAKIA R, et al. Cord blood metabolites associated with newborn adiposity and hyperinsulinemia. *J Pediatr*. 2018 Dec;203:144-149.e1. doi: 10.1016/j.jpeds.2018.07.056. Epub 2018 Sep 10. PMID: 30213459; PMCID: PMC6252151
24. SCHNEIDER C, et al. Associations of neonatal adiponectin and leptin with growth and body composition in African American infants. *Pediatr Obes*. 2018 Aug;13(8):485-491. doi: 10.1111/ijpo.12274. Epub 2018 Mar 24. PMID: 29573242; PMCID: PMC6457440
25. ANDERSEN G, et al. Body composition growth patterns in early infancy: A latent class trajectory analysis of the Ethiopian iABC Birth Cohort. *Obesity (Silver Spring)*. 2018 Jul;26(7):1225-1233. doi: 10.1002/oby.22197. Epub 2018 May 30. PMID: 29845756, 26(7):1225-1233 (2018)
26. ABERA M, et al. Body composition during early infancy and developmental progression from 1 to 5 years of age: The Infant Anthropometry and Body Composition (iABC) cohort study among Ethiopian children. *Br J Nutr*. 2018 Jun;119(11):1263-1273. doi: 10.1017/S000711451800082X. PMID: 29770755
27. DISSANAYAKE H, et al. Noninvasive assessment of autonomic function in human neonates born at the extremes of fetal growth spectrum. *Physiol Rep*. 2018 Apr;6(8):e13682. doi: 10.14814/phy2.13682. PMID: 29687617; PMCID: PMC5913591
28. ADMASSU B, et al. Accretion of fat-free mass rather than fat mass in infancy is positively associated with linear growth in childhood. *J Nutr*. 2018 Apr 1;148(4):607-615. doi: 10.1093/jn/nxy003. PMID: 29659955
29. DISSANAYAKE H, et al. Body Fatness and Cardiovascular Health in Newborn Infants, *J Clin Med*. 2018 Sep 11;7(9):270. doi: 10.3390/jcm7090270. PMID: 30208579; PMCID: PMC6162858
30. KADAKIA R, et al. Maternal pre-pregnancy BMI downregulates neonatal cord blood LEP methylation. *Pediatr Obes*. 2017 Aug;12 Suppl 1(Suppl 1):57-64. doi: 10.1111/ijpo.12204. Epub 2016 Dec 8. PMID: 27933755; PMCID: PMC5462869
31. ADMASSU B, et al. Body composition at birth and height at 2 years: A prospective cohort study among children in Jimma, Ethiopia. *Pediatr Res*. 2017 Aug;82(2):209-214. doi: 10.1038/pr.2017.59. Epub 2017 Jun 7. PMID: 28422940
32. SAUDER K, et al. Predictors of infant body composition at 5 months of age: The Healthy Start study. *J Pediatr*. 2017 Apr;183:94-99.e1. doi: 10.1016/j.jpeds.2017.01.014. Epub 2017 Feb 1. PMID: 28161200; PMCID: PMC5367947
33. BREIJ L, et al. Appetite-regulating hormones in early life and relationships with type of feeding and body composition in healthy term infants. *Eur J Nutr*. 2017 Jun;56(4):1725-1732. doi: 10.1007/s00394-016-1219-8. Epub 2016 May 11. PMID: 27170102; PMCID: PMC5486626
34. BREIJ L, et al. Longitudinal fat mass and visceral fat during the first 6 months after birth in healthy infants: Support for a critical window for adiposity in early life. *Pediatr Obes*. 2017 Aug;12(4):286-294. doi: 10.1111/ijpo.12139. Epub 2016 Apr 13. PMID: 27072083; PMCID: PMC6186414
35. CHAOIMH C, et al. Cord blood leptin and gains in body weight and fat mass during infancy. *Eur J Endocrinol*. 2016 Nov;175(5):403-10. doi: 10.1530/EJE-16-0431. Epub 2016 Aug 15. PMID: 27528502
36. PALEY C, et al. Body fat differences by self-reported race/ethnicity in healthy term newborns. *Pediatr Obes*. 2016 Oct;11(5):361-8. doi: 10.1111/ijpo.12072. Epub 2015 Oct 28. PMID: 26509351; PMCID: PMC4848178
37. KADAKIA R, et al. Neonatal adiposity increases with rising cord blood IGF-1 levels. *Clin Endocrinol (Oxf)*. 2016 Jul;85(1):70-5. doi: 10.1111/cen.13057. Epub 2016 Mar 31. PMID: 26945928; PMCID: PMC7036289
38. O'DONOVAN S, et al. Neonatal adiposity increases the risk of atopic dermatitis during the first year of life. *J Allergy Clin Immunol*. 2016 Jan;137(1):108-117. doi: 10.1016/j.jaci.2015.05.035. Epub 2015 Jul 17. PMID: 26194543
39. FAROOQ M, et al. Monitoring of infant feeding behavior using a jaw motion sensor. *J Healthc Eng*. 2015;6(1):23-40. doi: 10.1260/2040-2295.6.1.23. PMID: 25708375; PMCID: PMC4752004

40. O'DONOVAN S, et al. Cohort profile: The Cork BASELINE birth cohort study: Babies after SCOPE: Evaluating the longitudinal impact on neurological and nutritional endpoints. *Int J Epidemiol*. 2015 Jun;44(3):764-75. doi: 10.1093/ije/dyu157. Epub 2014 Aug 7. PMID: 25102856
41. HENRICKSSON P, et al. Variation in the fat mass and obesity-related (FTO) genotype is not associated with body fatness in infants, but possibly with their length. *Ped Obes*. 2014 Oct;9(5):e112-5. doi: 10.1111/ijpo.231. Epub 2014 May 21. PMID: 24846219
42. DONNELLEY E, et al. Antenatal predictors and body composition of large-for-gestational-age newborns: Perinatal health outcomes. *J Perinatol*. 2014 Sep;34(9):698-704. doi: 10.1038/jp.2014.90. Epub 2014 May 15. PMID: 24831524
43. GIANNI M, et al. Body composition changes in the first 6 months of life according to method of feeding. *J Hum Lact*. 2014 May;30(2):148-55. doi: 10.1177/0890334413516196. Epub 2013 Dec 18. PMID: 24352651
44. O'CONNOR C, et al. Birth weight and neonatal adiposity prediction using fractional limb volume obtained with 3D ultrasound. *Fetal Diag Ther*. 2014;36(1):44-8. doi: 10.1159/000360417. Epub 2014 Jun 6. PMID: 24924878
45. STANFIELD K, et al. Differences in body composition between infants of South Asian and European ancestry: the London Mother and Baby Study. *Int J Epidemiol*. 2012 Oct;41(5):1409-18. doi: 10.1093/ije/dys139. Epub 2012 Sep 14. PMID: 22984147; PMCID: PMC3465771
46. LEE W, et al. The relationship of newborn adiposity to fetal growth outcome based on birth weight or the modified neonatal growth assessment score. *J Matern Fetal Neonatal Med*. 2012 Oct;25(10):1933-40. doi: 10.3109/14767058.2012.683084. Epub 2012 May 29. PMID: 22494346; PMCID: PMC3930167
47. ERIKSSON B, et al. Body-composition development during early childhood and energy expenditure in response to physical activity in 1.5-year-old children. *Am J Clin Nutr*. 2012 Sep;96(3):567-73. doi: 10.3945/ajcn.111.022020. Epub 2012 Jul 25. PMID: 22836033
48. ANDERSEN G, et al. Fat and fat-free mass at birth: air displacement plethysmography measurements on 350 Ethiopian newborns. *Pediatr Res*. 2011 Nov;70(5):501-6. doi: 10.1203/PDR.0b013e31822d7470. PMID: 21772228
49. ERIKSSON B, et al. Fat-free mass hydration in newborns: Assessment and implications for body composition studies. *Acta Paediatr*. 2011 May;100(5):680-6. doi: 10.1111/j.1651-2227.2011.02147.x. Epub 2011 Feb 14. PMID: 21226759
50. ROGGERO P, et al. Neonatal period: Body composition changes in breast-fed full-term newborns. *Neonatology*. 2010;97(2):139-43. doi: 10.1159/000239767. Epub 2009 Sep 22. PMID: 19776647
51. ERIKSSON B, et al. Body composition in full-term healthy infants measured with air displacement plethysmography at 1 and 12 weeks of age. *Acta Paediatr*. 2010 Apr;99(4):563-8. doi: 10.1111/j.1651-2227.2009.01665.x. Epub 2010 Jan 8. PMID: 20064135
52. FIELDS D, et al. Sex differences in body composition early in life. *Gender Med*. 2009 Jul;6(2):369-75. doi: 10.1016/j.genm.2009.07.003. PMID: 19682664
53. MOYER-MILEUR L, et al. Newborn adiposity measured by plethysmography is not predicted by late gestation two-dimensional ultrasound measures of fetal growth. *J Nutr*. 2009 Sep;139(9):1772-8. doi: 10.3945/jn.109.109058. Epub 2009 Jul 29. PMID: 19640967; PMCID: PMC3151022

### **Mother/Newborn Relationship**

1. DEJANI N, et al. Maternal plasma transforming growth factor- $\beta$ 1 (TGF- $\beta$ 1) and newborn size: the Araraquara Cohort Study. *J Pediatr (Rio J)*. 2023 May-Jun;99(3):284-288. doi: 10.1016/j.jpeds.2022.11.009. Epub 2022 Dec 23. PMID: 36567066; PMCID: PMC10202726.
2. PRIORESCHI A, et al. Interactions between maternal health and placental morphology on neonate body composition. *Glob Pediatr*. 2023 Mar;3:100030. doi: 10.1016/j.gpeds.2022.100030. Epub 2022 Dec 5. PMID: 37073307; PMCID: PMC10107011.
3. MARTENIES S et al. Using non-parametric Bayes shrinkage to assess relationships between multiple environmental and social stressors and neonatal size and body composition in the Healthy Start cohort. *Environ Health*. 2022 Nov 19;21(1):111. doi: 10.1186/s12940-022-00934-z. PMID: 36401268; PMCID: PMC9675112.

4. HOYT A, et al. Prenatal exposure to tobacco and adverse birth outcomes: effect modification by folate intake during pregnancy. *Matern Health Neonatol Perinatol*. 2022 Sep 12;8(1):6. doi: 10.1186/s40748-022-00141-1. PMID: 36096906; PMCID: PMC9465971.
5. MOORE B, et al. Early-life exposure to tobacco and childhood adiposity: Identifying windows of susceptibility. *Pediatr Obes*. 2022 Dec;17(12):e12967. doi: 10.1111/ijpo.12967. Epub 2022 Aug 17. PMID: 36350199.
6. ASHWIN D, et al. The impact a Mediterranean Diet in the third trimester of pregnancy has on neonatal body fat percentage. *J Dev Orig Health Dis*. 2022 Aug;13(4):500-507. doi: 10.1017/S2040174421000556. Epub 2021 Oct 18.
7. FUCHS-NEUHOLD B, et al. Investigating New Sensory Methods Related to Taste Sensitivity, Preferences, and Diet of Mother-Infant Pairs and Their Relationship With Body Composition and Biomarkers: Protocol for an Explorative Study. *JMIR Res Protoc*. 2022 Apr 27;11(4):e37279. doi: 10.2196/37279. PMID: 35475790 PMCID: PMC9096638
8. WHYTE K, et al. A Secondary Analysis of Maternal Ultra-Processed Food Intake in Women with Overweight or Obesity and Associations with Gestational Weight Gain and Neonatal Body Composition Outcomes. *J Mother Child*. 2022 Mar 23. doi: 10.34763/jmotherandchild.20212504.d-21-00025. eCollection 2021 Dec 1. PMID: 35325513 PMCID: PMC9444195
9. NAKANDAKARE P, et al. Associations between obesity-related gene expression in maternal and cord blood and newborn adiposity: Findings from the Araraquara Cohort study. *Int J Obes (Lond)*. 2021 Sep;45(9):1958-1966. doi: 10.1038/s41366-021-00857-8. Epub 2021 May 17. PMID: 34002037
10. WHYTE K, et al. No sustained effects of an intervention to prevent excessive GWG on offspring fat and lean mass at 54 weeks: Yet a greater head circumference persists. *Pediatr Obes*. 2021 Jul;16(7):e12767. doi: 10.1111/ijpo.12767. Epub 2021 Jan 4. PMID: 33394566; PMCID: PMC8178185
11. HERATH MP, et al. Determinants of Infant Adiposity across the First 6 Months of Life: Evidence from the Baby-Bod study. *J Clin Med*. 2021 Apr 19;10(8):1770. doi: 10.3390/jcm10081770. PMID: 33921680; PMCID: PMC8073882
12. WIECHERS C, et al. Body composition in term offspring after maternal gestational diabetes does not predict postnatal hypoglycemia. *BMC Pediatr*. 2021 Mar 6;21(1):111. doi: 10.1186/s12887-021-02578-3. PMID: 33676430; PMCID: PMC7936473
13. WOLFS D, et al. Brown fat-activating lipokine 12,13-diHOME in human milk is associated with infant adiposity. *J Clin Endocrinol Metab*. 2021 Jan 23;106(2):e943-e956. doi: 10.1210/clinem/dgaa799. PMID: 33135728; PMCID: PMC7823229
14. NEHAB S, et al. Influence of gestational and perinatal factors on body composition of full-term newborns. *J Pediatr (Rio J)*. 2020 Nov-Dec;96(6):771-777. doi: 10.1016/j.jped.2019.09.006. Epub 2019 Nov 9. PMID: 31711787
15. MOORE B, et al. Neonatal adiposity and childhood obesity. *Pediatrics*. 2020 Sep;146(3):e20200737. doi: 10.1542/peds.2020-0737. Epub 2020 Aug 13. PMID: 32796097; PMCID: PMC7461209
16. MATUSIAK K, et al. Self-reported periconception weight loss attempts do not alter infant body composition. *Nutrition*. 2020 Sep;77:110781. doi: 10.1016/j.nut.2020.110781. Epub 2020 Feb 29. PMID: 32283340
17. ANDERSSON-HALL U, et al. Infant body composition relationship to maternal adipokines and fat mass: The PONCH study. *Pediatr Res*. 2020 Sep 14. doi: 10.1038/s41390-020-01115-9. Epub ahead of print. PMID: 3292747
18. WROTTESELEY S, et al. The association between fetal growth and neonatal adiposity in urban South African infants. *Pediatr Obes*. 2020 Sep 30:e12737. doi: 10.1111/ijpo.12737. Epub ahead of print. PMID: 32997439
19. NEHAB S, et al. Gestational weight gain and body composition of full-term newborns and infants: A cohort study. *BMC Pregnancy Childbirth*. 2020 Aug 20;20(1):474. doi: 10.1186/s12884-020-03145-x. PMID: 32819310; PMCID: PMC7439530
20. ERICKSON M, et al. Rationale and study design for lifestyle intervention in preparation for pregnancy (LIPP): A randomized controlled trial. *Contemp Clin Trials*. 2020 Jul;94:106024. doi: 10.1016/j.cct.2020.106024. Epub 2020 May 8. PMID: 32389808; PMCID: PMC7375196
21. DIAZ E, et al. Parental adiposity differentially associates with newborn body composition. *Pediatr Obes*. 2020 Apr;15(4):e12596. doi: 10.1111/ijpo.12596. Epub 2019 Dec 19. PMID: 31856430; PMCID: PMC7367307
22. STARLING A, et al. Prenatal exposure to traffic and ambient air pollution and infant weight and adiposity: The Healthy Start study. *Environ Res*. 2020 Mar;182:109130. doi: 10.1016/j.envres.2020.109130. Epub 2020 Jan 10. PMID: 32069764; PMCID: PMC7394733

23. DISSANAYAKE H, et al. Influence of maternal and placental factors on newborn body composition. *J Paediatr Child Health*. 2020 Feb;56(2):224-230. doi: 10.1111/jpc.14565. Epub 2019 Aug 7. PMID: 31392795
24. BENNETT A, et al. Maternal sociodemographic and health behaviours associated with adiposity in infants as measured by air displacement plethysmography. *Early Hum Dev*. 2019 Oct 26;140:104887. doi: 10.1016/j.earlhumdev.2019.104887. Epub ahead of print. PMID: 31670177
25. HOLLANDERS J, et al. No association between glucocorticoid diurnal rhythm in breastmilk and infant body composition at 3 months. *Nutrients*. 2019 Oct 2;11(10):2351. doi: 10.3390/nu11102351. PMID: 31581748; PMCID: PMC6835896
26. ABREU L, et al. Gestational diabetes mellitus, pre-pregnancy body mass index, and gestational weight gain as risk factors for increased fat mass in Brazilian newborns. *PLoS One*. 2019 Aug 29;14(8):e0221971. doi: 10.1371/journal.pone.0221971. PMID: 31465493; PMCID: PMC6715169
27. VAN LEE L, et al. Prospective associations of maternal choline status with offspring body composition in the first 5 years of life in two large mother-offspring cohorts: The Southampton Women's Survey cohort and the Growing Up in Singapore Towards healthy Outcomes cohort. *Int J Epidemiol*. 2019 Apr 1;48(2):433-444. doi: 10.1093/ije/dyy291. PMID: 30649331; PMCID: PMC6751083
28. MUDD L, et al. Relations among maternal physical activity during pregnancy and child body composition. *Obes Sci Pract*. 2019 Mar 1;5(3):246-250. doi: 10.1002/osp4.332. PMID: 31275598; PMCID: PMC6587308
29. FUENTES E, et al. Parental obesity differentially associates with newborn adiposity. Is programming of the GH-IGF axis involved? *Curr Dev Nutr*. 2019;3(Suppl 1):nzz041.OR09-05-19. Published 2019 Jun 13. doi:10.1093/cdn/nzz041.OR09-05-19
30. MOORE B, et al. Fetal exposure to maternal active and secondhand smoking with offspring early-life growth in the Healthy Start study. *Int J Obes*. 2019 Apr;43(4):652-662. doi: 10.1038/s41366-018-0238-3. Epub 2018 Oct 19. PMID: 30341407; PMCID: PMC6445676
31. ANDERSSON-HALL U, et al. Maternal obesity and gestational diabetes mellitus affect body composition through infancy: The PONCH study. *Pediatr Res*. 2019 Feb;85(3):369-377. doi: 10.1038/s41390-018-0248-9. Epub 2018 Dec 10. PMID: 30705398
32. KENNEDY R, et al. The relationship between early pregnancy dietary intakes and subsequent birthweight and neonatal adiposity. *J Pub Health*. 2018 Dec 1;40(4):747-755. doi: 10.1093/pubmed/fdx131. PMID: 30590769
33. PHANG M, et al. Increased  $\alpha$ -linolenic acid intake during pregnancy is associated with higher offspring birth weight. *Curr Dev Nutr*. 2018 Oct 18;3(2):nzy081. doi: 10.1093/cdn/nzy081. PMID: 30820488; PMCID: PMC6390041
34. MACAULAY S, et al. The effects of gestational diabetes mellitus on fetal growth and neonatal birth measures in an African cohort. *Diabet Med*. 2018 Oct;35(10):1425-1433. doi: 10.1111/dme.13668. Epub 2018 Jun 1. PMID: 29766563
35. WROTTESELEY S, et al. Maternal traditional dietary pattern and antiretroviral treatment exposure are associated with neonatal size and adiposity in urban, black South Africans. *Br J Nutr*. 2018 Sep;120(5):557-566. doi: 10.1017/S0007114518001708. Epub 2018 Jul 30. PMID: 30058507; PMCID: PMC6773599
36. BARBOUR L, et al. Postprandial triglycerides predict newborn fat more strongly than glucose in women with obesity in early pregnancy. *Obesity*. 2018 Aug;26(8):1347-1356. doi: 10.1002/oby.22246. Epub 2018 Jun 22. PMID: 29931812; PMCID: PMC6107410
37. OKESENE-GAFA K, et al. A randomised controlled demonstration trial of multifaceted nutritional intervention and or probiotics: the healthy mums and babies (HUMBA) trial. *BMC Pregnancy Childbirth*. 2018 May 4;18(1):130. PMID: 27884128; PMCID: PMC5123375
38. MOORE B, et al. Proinflammatory diets during pregnancy and neonatal adiposity in the Healthy Start Study. *J Pediatr*. 2018 Apr;195:121-127.e2. doi: 10.1016/j.jpeds.2017.10.030. Epub 2017 Dec 6. PMID: 29217099; PMCID: PMC6363107
39. GALLAGHER D, et al. Greater neonatal fat-free mass and similar fat mass following a randomized trial to control excess gestational weight gain. *Obesity*. 2018 Mar;26(3):578-587. doi: 10.1002/oby.22079. PMID: 29464905; PMCID: PMC5824435
40. DAHLY D, et al. (Cork) cohort study Associations between maternal lifestyle factors and neonatal body composition in the Screening for Pregnancy Endpoints. *Int J Epidemiol*. 2018 Feb 1;47(1):131-145. doi: 10.1093/ije/dyx221. PMID: 29136159
41. DAHLY D, et al. Associations between maternal lifestyle factors and neonatal body composition in the Screening for Pregnancy Endpoints (Cork) cohort study. *Int J Epidemiol*. 2018 Feb 1;47(1):131-145. doi: 10.1093/ije/dyx221. PMID: 29136159

42. CHIA A, et al. Adherence to a healthy eating index for pregnant women is associated with lower neonatal adiposity in a multiethnic Asian cohort: The Growing Up in Singapore Towards Healthy Outcomes (GUSTO) Study. *Am J Clin Nutr.* 2018 Jan 1;107(1):71-79. doi: 10.1093/ajcn/nqx003. PMID: 29381790; PMCID: PMC5972656
43. NORRIS T, et al. Do changing levels of maternal exercise during pregnancy affect neonatal adiposity? Secondary analysis of the babies after SCOPE: evaluating the longitudinal impact using neurological and nutritional endpoints (BASELINE) birth cohort (Cork, Ireland). *BMJ Open.* 2017 Dec 1;7(11):e017987. doi: 10.1136/bmjopen-2017-017987. PMID: 29196482; PMCID: PMC5719326
44. MCKENZIE K, et al. Quantity and quality of carbohydrate intake during pregnancy, newborn body fatness and cardiac autonomic control: Conferred cardiovascular risk? *Nutrients.* 2017 Dec 19;9(12):1375. doi: 10.3390/nu9121375. PMID: 29257088; PMCID: PMC5748825
45. BAKER P 2<sup>nd</sup>, et al. Altered gene expression and metabolism in fetal umbilical cord mesenchymal stem cells correspond with differences in 5-month-old infant adiposity gain. *Sci Rep.* 2017 Dec 22;7(1):18095. doi: 10.1038/s41598-017-17588-4. PMID: 29273781; PMCID: PMC5741772
46. BAKER P 2<sup>nd</sup>, et al. Maternal obesity and increased neonatal adiposity correspond with altered infant mesenchymal stem cell metabolism. *JCI Insight.* 2017 Nov 2;2(21):e94200. doi: 10.1172/jci.insight.94200. PMID: 29093265; PMCID: PMC5752288
47. BREIJ L, et al. Impact of early infant growth, duration of breastfeeding and maternal factors on total body fat mass and visceral fat at 3 and 6 months of age. *Ann Nutr Metab.* 2017;71(3-4):203-210. doi: 10.1159/000481539. Epub 2017 Nov 14. PMID: 29136614; PMCID: PMC6198935
48. MACDONALD T, et al. Reduced growth velocity across the third trimester is associated with placental insufficiency in fetuses born at a normal birthweight: A prospective cohort study. *BMC Med.* 2017 Aug 31;15(1):164. doi: 10.1186/s12916-017-0928-z. PMID: 28854913; PMCID: PMC5577811.
49. FUGLESTAD A, et al. Maternal executive function, infant feeding responsiveness and infant growth during the first 3 months. *Pediatr Obes.* 2017 Aug;12 Suppl 1:102-110. doi: 10.1111/ijpo.12226. Epub 2017 Jul 28. PMID: 28752657
50. MOORE B, et al. Exposure to secondhand smoke, exclusive breastfeeding and infant adiposity at age 5 months in the Healthy Start study. *Pediatr Obes.* 2017 Aug;12 Suppl 1(Suppl 1):111-119. doi: 10.1111/ijpo.12233. PMID: 28868814; PMCID: PMC5608262
51. SAUDER K, et al. Prenatal vitamin D intake, cord blood 25-hydroxyvitamin D, and offspring body composition: The Healthy Start Study. *Nutrients.* 2017 Jul 22;9(7):790. doi: 10.3390/nu9070790. PMID: 28737667; PMCID: PMC5537904
52. BERNARD J, et al. Maternal plasma phosphatidylcholine polyunsaturated fatty acids during pregnancy and offspring growth and adiposity. *Prostaglandins Leukot Essent Fatty Acids.* 2017 Jun;121:21-29. doi: 10.1016/j.plefa.2017.05.006. Epub 2017 Jun 9. PMID: 28651694; PMCID: PMC5501311
53. STARLING A, et al. Perfluoroalkyl substances during pregnancy and offspring weight and adiposity at birth: Examining mediation by maternal fasting glucose in the Healthy Start Study. *Environ Health Perspect.* 2017 Jun 26;125(6):067016. doi: 10.1289/EHP641. PMID: 28669937; PMCID: PMC5743451
54. RUDOLPH M, et al. Early infant adipose deposition is positively associated with the n-6 to n-3 fatty acid ratio in human milk independent of maternal BMI. *Int J Obes (Lond).* 2017 Apr;41(4):510-517. doi: 10.1038/ijo.2016.211. Epub 2016 Nov 23. PMID: 27876761; PMCID: PMC5380514
55. CASTRO N, et al. The relationship between maternal plasma leptin and adiponectin concentrations and newborn adiposity. *Nutrients.* 2017 Feb 23;9(3):182. doi: 10.3390/nu9030182. PMID: 28241462; PMCID: PMC5372845
56. LOGAN K, et al. Diabetes in pregnancy and infant adiposity: systematic review and meta-analysis. *Arch Dis Child Fetal Neonatal Ed.* 2017 Jan;102(1):F65-F72. doi: 10.1136/archdischild-2015-309750. Epub 2016 May 26. PMID: 27231266; PMCID: PMC5256410
57. JOSEFSON J, et al. Excessive gestational weight gain in the first trimester among women with normal glucose tolerance and resulting neonatal adiposity. *J Perinatol.* 2016 Dec;36(12):1034-1038. doi: 10.1038/jp.2016.145. Epub 2016 Sep 1. PMID: 27583397; PMCID: PMC5130601

58. SIMONI J, et al. A comparison of intestinal microbiota in a population of low-risk infants exposed and not exposed to intrapartum antibiotics: The Baby & Microbiota of the Intestine cohort study protocol. *BMC Pediatr.* 2016 Nov 10;16(1):183. doi: 10.1186/s12887-016-0724-5. PMID: 27832763; PMCID: PMC510339
59. CRUME T, et al. Maternal dietary intake during pregnancy and offspring body composition: The Healthy Start Study. *Am J Obstet Gynecol.* 2016 Nov;215(5):609.e1-609.e8. doi: 10.1016/j.ajog.2016.06.035. Epub 2016 Jun 28. PMID: 27371352; PMCID: PMC5571832
60. PEREIRA-DA-SILVA L, et al. Low adherence to Mediterranean diet in Portugal: Pregnant women nutrition in Portugal and its repercussions. *Acta Med Port.* 2016 Oct;29(10):658-666. doi: 10.20344/amp.7344. Epub 2016 Oct 31. PMID: 28103462
61. MCCARTHY F, et al. Parental physical and lifestyle factors and their association with newborn body composition. *BJOG.* 2016 Oct;123(11):1824-9. doi: 10.1111/1471-0528.14042. Epub 2016 Apr 22. PMID: 27102226
62. SAUDER K, et al. Exploring the association between maternal prenatal multivitamin use and early infant growth: The Healthy Start Study. *Pediatr Obes.* 2016 Oct;11(5):434-41. doi: 10.1111/ijpo.12084. Epub 2015 Dec 11. PMID: 26663829; PMCID: PMC4903091
63. SHAPIRO A, et al. Nicotinamide promotes adipogenesis in umbilical cord-derived mesenchymal stem cells and is associated with neonatal adiposity: The Healthy Start BabyBUMP Project. *PLoS One.* 2016 Jul 14;11(7):e0159575. doi: 10.1371/journal.pone.0159575. PMID: 27414406; PMCID: PMC4944979
64. SHAPIRO A, et al. Maternal diet quality in pregnancy and neonatal adiposity: The Healthy Start Study. *Int J Obes.* 2016 Jul;40(7):1056-62. doi: 10.1038/ijo.2016.79. Epub 2016 May 2. PMID: 27133623; PMCID: PMC5356926
65. LEMAS D, et al. Alterations in human milk leptin and insulin are associated with early changes in the infant intestinal microbiome. *Am J Clin Nutr.* 2016 May;103(5):1291-300. doi: 10.3945/ajcn.115.126375. PMID: 27140533; PMCID: PMC4841936
66. KIZIRIAN N, et al. Macronutrient balance and dietary glycemic index in pregnancy predict neonatal body composition. *Nutrients.* 2016 May 6;8(5):270. doi: 10.3390/nu8050270. PMID: 27164136; PMCID: PMC4882683
67. KIZIRIAN N, et al. Effects of a low-glycemic index diet during pregnancy on offspring growth, body composition, and vascular health: A pilot randomized controlled trial. *Am J Clin Nutr.* 2016 Apr;103(4):1073-82. doi: 10.3945/ajcn.115.123695. Epub 2016 Mar 2. PMID: 26936333
68. TINIUS R, et al. Maternal inflammation during late pregnancy is lower in physically active compared with inactive obese women. *Appl Physiol Nutr Metab.* 2016 Feb;41(2):191-8. doi: 10.1139/apnm-2015-0316. Epub 2015 Nov 3. PMID: 26799789; PMCID: PMC4883098
69. MARKOVIC T, et al. Randomized controlled trial investigating the effects of a low-glycemic index diet on pregnancy outcomes in women at high risk of gestational diabetes mellitus: the GI Baby 3 Study. *Diabetes Care.* 2016 Jan;39(1):31-8. doi: 10.2337/dc15-0572. Epub 2015 Jul 16. PMID: 26185283
70. LEMAS D, et al. Associations of maternal weight status prior and during pregnancy with neonatal cardiometabolic markers at birth: The Healthy Start Study. *Int J Obes (Lond).* 2015 Oct;39(10):1437-42. doi: 10.1038/ijo.2015.109. Epub 2015 Jun 9. PMID: 26055075; PMCID: PMC4596750
71. HENRIKSSON P, et al. Gestational weight gain according to Institute of Medicine recommendations in relation to infant size and body composition. *Pediatr Obes.* 2015 Oct;10(5):388-94. doi: 10.1111/ijpo.276. Epub 2014 Dec 17. PMID: 25521831
72. HULL H, et al. Maternal obesity influences the relationship between location of neonate fat mass and total fat mass. *Pediatr Obes.* 2015 Aug;10(4):245-51. doi: 10.1111/ijpo.257. Epub 2014 Aug 4. PMID: 25088238; PMCID: PMC431739
73. TINIUS R, et al. Altered maternal lipid metabolism is associated with higher inflammation in obese women during late pregnancy. *Integr Obes Diabetes.* 2015;2(1):168-175. doi: 10.15761/iob.1000137. Epub 2015 Dec 19. PMID: 27239331; PMCID: PMC4883583
74. BREIJ L, et al. Maternal and fetal determinants of neonatal body composition. *Horm Res Paediatr.* 2015;84(6):388-95. doi: 10.1159/000441298. Epub 2015 Oct 23. PMID: 26492188
75. HENRIKSSON P, et al. Glucose homeostasis variables in pregnancy versus maternal and infant body composition. *Nutrients.* 2015 Jul 10;7(7):5615-27. doi: 10.3390/nu7075243. PMID: 26184296; PMCID: PMC4517020

76. HENRIKSSON P, et al. Parental fat-free mass is related to the fat-free mass of infants and maternal fat mass is related to the fat mass of infant girls. *Acta Paediatr.* 2015 May;104(5):491-7. doi: 10.1111/apa.12939. Epub 2015 Mar 21. PMID: 2564582
77. CRUME T, et al. Maternal fuels and metabolic measures during pregnancy and neonatal body composition: The Healthy Start Study. *J Clin Endocrinol Metab.* 2015 Apr;100(4):1672-80. doi: 10.1210/jc.2014-2949. Epub 2015 Jan 9. PMID: 25574704; PMCID: PMC4399301
78. STARLING A, et al. Associations of maternal BMI and gestational weight gain with neonatal adiposity in The Healthy Start Study. *Am J Clin Nutr.* 2015 Feb;101(2):302-9. doi: 10.3945/ajcn.114.094946. Epub 2014 Dec 3. PMID: 25646327; PMCID: PMC4307203
79. HARROD C, et al. Exposure to prenatal smoking and early-life body composition: The Healthy Start Study. *Obesity.* 2015 Jan;23(1):234-41. doi: 10.1002/oby.20924. Epub 2014 Nov 11. PMID: 25385660; PMCID: PMC4276469
80. PEREIRA-DA-SILVA L, et al. The adjusted effect of body mass index, energy and macronutrient intakes during pregnancy, and gestational weight gain on body composition of full-term neonates. *Am J Perinatol.* 2014 Nov;31(10):875-82. doi: 10.1055/s-0033-1363502. Epub 2013 Dec 17. PMID: 24347258
81. HARROD C, et al. Quantity and timing of maternal prenatal smoking on neonatal body composition: The Healthy Start Study. *J Pediatr.* 2014 Oct;165(4):707-12. doi: 10.1016/j.jpeds.2014.06.031. Epub 2014 Jul 22. PMID: 25063722; PMCID: PMC4177331
82. HARROD C, et al. Physical activity in pregnancy and neonatal body composition: The Healthy Start Study. *Obstet Gynecol.* 2014 Aug;124(2 Pt 1):257-264. doi: 10.1097/AOG.0000000000000373. PMID: 25004346; PMCID: PMC4111108
83. ESTAMPADOR A, et al. Infant body composition and adipokine concentrations in relation to maternal gestational weight gain. *Diabetes Care.* 2014 May;37(5):1432-8. doi: 10.2337/dc13-2265. Epub 2014 Mar 12. PMID: 24623025; PMCID: PMC3994936
84. JOSEFSON J, et al. Maternal leptin predicts adiposity of the neonate. *Horm Res Paediatr.* 2014;81(1):13-9. doi: 10.1159/000355387. Epub 2013 Dec 11. PMID: 24334975; PMCID: PMC4123455
85. CARBERRY A, et al. Customized versus population-based birth weight charts for the detection of neonatal growth and perinatal morbidity in a cross-sectional study of term neonates. *Am J Epidemiol.* 2013 Oct 15;178(8):1301-8. doi: 10.1093/aje/kwt176. Epub 2013 Aug 21. PMID: 23966560
86. AU C, et al. Fetal and maternal factors associated with neonatal adiposity as measured by air displacement plethysmography: a large cross-sectional study. *Early Hum Dev.* 2013 Oct;89(10):839-43. doi: 10.1016/j.earlhumdev.2013.07.028. Epub 2013 Aug 19. PMID: 23968962
87. JOSEFSON J, et al. Excessive weight gain in women with a normal pre-pregnancy BMI is associated with increased neonatal adiposity. *Ped Obes.* 2013 Apr;8(2):e33-6. doi: 10.1111/j.2047-6310.2012.00132.x. Epub 2013 Jan 3. PMID: 23283756; PMCID: PMC4076951
88. AU C, et al. Body composition is normal in term infants born to mothers with well-controlled gestational diabetes mellitus. *Diabetes Care.* 2013 Mar;36(3):562-4. doi: 10.2337/dc12-1557. Epub 2012 Dec 6. PMID: 23223404; PMCID: PMC3579380
89. POMEROY J, et al. Maternal physical activity and insulin action in pregnancy and their relationships with infant body composition. *Diabetes Care.* 2013 Feb;36(2):267-9. doi: 10.2337/dc12-0885. Epub 2012 Sep 10. PMID: 22966095; PMCID: PMC3554315
90. JOSEFSON J, et al. Maternal obesity and vitamin D sufficiency are associated with cord blood vitamin D insufficiency. *J Clin Endocrinol Metab.* 2013 Jan;98(1):114-9. doi: 10.1210/jc.2012-2882. Epub 2012 Nov 8. PMID: 23144468; PMCID: PMC3537089
91. LARSON ODE K, et al. Decelerated early growth in infants of overweight and obese mothers. *J Pediatr.* 2012 Dec;161(6):1028-34. doi: 10.1016/j.jpeds.2012.06.001. Epub 2012 Jul 20. PMID: 22819273; PMCID: PMC3480982
92. ANDRES A. et al. Body fat mass of exclusively breastfed infants born to overweight mothers. *J Acad Nutr Diet.* 2012 Jul;112(7):991-5. doi: 10.1016/j.jand.2012.03.031. PMID: 22889630
93. HULL H, et al. Higher infant body fat with excessive gestational weight gain in overweight women. *Am J Obstet Gynecol.* 2011 Sep;205(3):211.e1-7. doi: 10.1016/j.ajog.2011.04.004. Epub 2011 Apr 14. PMID: 21621185; PMCID: PMC3170486
94. HULL H, et al. Impact of maternal body mass index on neonate birthweight and body composition. *Am J Obstet Gynecol.* 2008 Apr;198(4):416.e1-6. doi: 10.1016/j.ajog.2007.10.796. Epub 2008 Feb 15. PMID: 18279830

## Body Composition and Nutritional Intervention

1. CARDOSO M, et al. Individualized Fortification Based on Measured Macronutrient Content of Human Milk Improves Growth and Body Composition in Infants Born Less than 33 Weeks: A Mixed-Cohort Study. *Nutrients*. 2023 Mar 22;15(6):1533. doi: 10.3390/nu15061533. PMID: 36986263; PMCID: PMC10052754.
2. MARDERFELD L, et al. Body composition correlates with laboratory parameters and disease severity in infants with biliary atresia. *Pediatr Transplant*. 2022 May;26(3):e14208. doi: 10.1111/ptr.14208. Epub 2021 Dec 19. PMID: 34927330
3. VLUG LE, Anthropometrics and fat mass, but not fat-free mass, are compromised in infants requiring parenteral nutrition after neonatal intestinal surgery. *Am J Clin Nutr*. 2022 Feb 9;115(2):503-513. doi: 10.1093/ajcn/nqab345. PMID: 34637493 PMCID: PMC8827070
4. WIECHERS C, et al. Body Composition of Preterm Infants following Rapid Transition to Enteral Feeding. *Neonatology*. 2022;119(2):246-254. doi: 10.1159/000521032. Epub 2022 Jan 17. PMID: 35038714
5. ALJA'NINI Z, et al. Effect of parenteral nutrition duration on patterns of growth and body composition in very low birth weight premature infants. *JPEN J Parenter Enteral Nutr*. 2021 Nov;45(8):1673-1682. doi: 10.1002/jpen.2278. Epub 2021 Oct 29. PMID: 34638161
6. OLGA L, Extensive Study of Breast Milk and Infant Growth: Protocol of the Cambridge Baby Growth and Breastfeeding Study (CBGS-BF). *Nutrients*. 2021 Aug 21;13(8):2879. doi: 10.3390/nu13082879. PMID: 34445039; PMCID: PMC8400677
7. ALJA'NINI Z, et al. Infant body composition assessment in the neonatal intensive care unit (NICU) using air displacement plethysmography: Strategies for implementation into clinical workflow. *Clin Nutr ESPEN*. 2021 Jun;43:212-222. doi: 10.1016/j.clnesp.2021.04.014. Epub 2021 Apr 24. PMID: 34024517
8. KOUWENHOVEN SMP, et al. Long-term effects of a modified, low-protein infant formula on growth and body composition: Follow-up of a randomized, double-blind, equivalence trial. *Clin Nutr*. 2021 Jun;40(6):3914-3921. doi: 10.1016/j.clnu.2021.04.034. Epub 2021 Apr 27. PMID: 34139464
9. OLHAGER E, et al. Different feeding regimens were not associated with variation in body composition in preterm infants. *J Matern Fetal Neonatal Med*. 2021 Jun 7:1-8. doi: 10.1080/14767058.2021.1914575. Epub ahead of print. PMID: 34098845
10. JAYASINGHE S, et al. Exclusivity of breastfeeding and body composition: Learnings from the Baby-Bod Study. *Int Breastfeed J*. 2021 May 19;16(1):41. doi: 10.1186/s13006-021-00389-x. PMID: 34011366; PMCID: PMC8132405
11. VISWANATHAN S, et al. Body adiposity and oral feeding outcomes in infants: A pilot study. *J Perinatol*. 2021 May;41(5):1059-1064. doi: 10.1038/s41372-021-00975-1. Epub 2021 Feb 17. PMID: 33597738
12. BELFORT MB, et al. Targeting human milk fortification to improve very preterm infant growth and brain development: study protocol for Nourish, a single-center randomized, controlled clinical trial. *BMC Pediatr*. 2021 Apr 9;21(1):167. doi: 10.1186/s12887-021-02635-x. PMID: 33836708; PMCID: PMC8033746
13. SALAS A, et al. Percent body fat content measured by plethysmography in infants randomized to high- or usual-volume feeding after very preterm birth. *J Pediatr*. 2021 Mar;230:251-254.e3. doi: 10.1016/j.jpeds.2020.11.028. Epub 2020 Nov 25. PMID: 33248115; PMCID: PMC7914146
14. CARDOSO M, et al. Customized human milk fortification based on measured human milk composition to Improve the quality of growth in very preterm infants: A mixed-cohort study protocol. *Int J Environ Res Public Health*. 2021 Jan 19;18(2):823. doi: 10.3390/ijerph18020823. PMID: 33477964; PMCID: PMC7835734
15. ROCHOW N, et al. Individualized target fortification of breast milk with protein, carbohydrates, and fat for preterm infants: A double-blind randomized controlled trial. *Clin Nutr*. 2021 Jan;40(1):54-63. doi: 10.1016/j.clnu.2020.04.031. Epub 2020 May 6. PMID: 32446787
16. NAGEL E, et al. Clinical application of body composition methods in premature infants. *J Parenter Enteral Nutr*. 2020 Jul;44(5):785-795. doi: 10.1002/jpen.1803. Epub 2020 Feb 6. PMID: 32026515
17. MC NELIS K, et al. Body composition of very low-birth-weight infants fed fortified human milk: A pilot study. *J Parenter Enteral Nutr*. 2020 May 26. doi: 10.1002/jpen.1931. Epub ahead of print. PMID: 32458470

18. LIOTTO N, et al. Protein intakes during weaning from parenteral nutrition drive growth gain and body composition in very low birth weight preterm infants. *Nutrients*. 2020 May 2;12(5):1298. doi: 10.3390/nu12051298. PMID: 32370158; PMCID: PMC7282247
19. PARAT S, et al. Targeted Breast Milk Fortification for Very Low Birth Weight (VLBW) Infants: Nutritional Intake, Growth Outcome and Body Composition. *Nutrients*. 2020 Apr 21;12(4):1156. doi: 10.3390/nu12041156. PMID: 32326177 PMCID: PMC7230830
20. LINGWOOD B, et al. Body composition in very preterm infants before discharge is associated with macronutrient intake. *Br J Nutr*. 2020 Apr 14;123(7):800-806. doi: 10.1017/S000711451900343X. Epub 2020 Jan 6. PMID: 31902375
21. ATCHLEY C, et al. Enhanced protein diet for preterm infants: A prospective, randomized, double-blind, controlled trial. *J Pediatr Gastroenterol Nutr*. 2019 Aug;69(2):218-223. doi: 10.1097/MPG.0000000000002376. PMID: 31058772
22. MACEDO I, et al. Associations of measured protein and energy intakes with growth and adiposity in human milk-fed preterm infants at term postmenstrual age: A cohort study. *Am J Perinatol*. 2018 Jul;35(9):882-891. doi: 10.1055/s-0038-1626717. Epub 2018 Feb 2. PMID: 29532450
23. MORLACCHI L, et al. Protein use and weight-gain quality in very-low-birth-weight preterm infants fed human milk or formula. *Am J Clin Nutr*. 2018 Feb 1;107(2):195-200. doi: 10.1093/ajcn/nqx001. PMID: 29529139
24. LIOTTO N, Clinical evaluation of two different protein content formulas fed to full-term healthy infants: A randomized controlled trial. *BMC Pediatr*. 2018 Feb 13;18(1):59. doi: 10.1186/s12887-018-1046-6. Erratum in: *BMC Pediatr*. 2018 Apr 12;18(1):135. PMID: 29439736; PMCID: PMC5812222
25. GIANNI M, et al. Does human milk modulate body composition in late preterm infants at term-corrected age? *Nutrients*. 2016 Oct 23;8(10):664. doi: 10.3390/nu8100664. PMID: 27782098; PMCID: PMC5084050
26. TELLER I, et al. Post-discharge formula feeding in preterm infants: A systematic review mapping evidence about the role of macronutrient enrichment. *Clin Nutr*. 2016 Aug;35(4):791-801. doi: 10.1016/j.clnu.2015.08.006. Epub 2015 Sep 5. PMID: 26499034
27. BRENNAN A, et al. Optimising preterm nutrition: present and future. *Proc Nutr Soc*. 2016 May;75(2):154-61. doi: 10.1017/S0029665116000136. Epub 2016 Apr 1. PMID: 27032990
28. MCCLEOD G, et al. Comparing different methods of human breast milk fortification using measured v. assumed macronutrient composition to target reference growth: A randomised controlled trial. *Br J Nutr*. 2016 Feb 14;115(3):431-9. doi: 10.1017/S0007114515004614. Epub 2015 Dec 2. PMID: 26627899
29. MCLEOD G, et al. Feasibility study: Assessing the influence of macronutrient intakes on preterm body composition, using air displacement plethysmography. *J Paediatr Child Health*. 2015 Sep;51(9):862-9. doi: 10.1111/jpc.12893. Epub 2015 Apr 14. PMID: 25873446
30. BLOOMFIELD F, et al. The ProVIDE study: The impact of protein intravenous nutrition on development in extremely low birthweight babies. *BMC Pediatr*. 2015 Aug 26;15:100. doi: 10.1186/s12887-015-0411-y. PMID: 26307566; PMCID: PMC4549896
31. PEREIRA-DA-SILVA L, et al. Resting energy expenditure, macronutrient utilization, and body composition in term infants after corrective surgery of major congenital anomalies: A case-study. *J Neonatal Perinatal Med*. 2015;8(4):403-12. doi: 10.3233/NPM-15915019. PMID: 26757004
32. PEREIRA-DA-SILVA L, et al. The effect of long-chain polyunsaturated fatty acids intake during pregnancy on adiposity of healthy full-term offspring at birth. *J Perinatol*. 2015 Mar;35(3):177-80. doi: 10.1038/jp.2014.188. Epub 2014 Oct 16. PMID: 25321648
33. GIANNI M, et al. Formula-fed infants have significantly higher fat-free mass content in their bodies than breastfed babies. *Acta Paediatr*. 2014 Jul;103(7):e277-81. doi: 10.1111/apa.12643. Epub 2014 Apr 17. PMID: 24673117
34. TIMBY N, et al. Neurodevelopment, nutrition, and growth until 12 mo. of age in infants fed a low-energy, low-protein formula supplemented with bovine milk fat globule membranes: A randomized controlled trial. *Am J Clin Nutr*. 2014 Apr;99(4):860-8. doi: 10.3945/ajcn.113.064295. Epub 2014 Feb 5. PMID: 24500150
35. ROGGERO P, et al. Growth and fat-free mass gain in preterm infants after discharge: A randomized controlled trial. *Pediatrics*. 2012 Nov;130(5):e1215-21. doi: 10.1542/peds.2012-1193. Epub 2012 Oct 29. PMID: 23109680

- 36. ROGGERO P, et al. Effect of nutrition on growth and body composition in infants born preterm. *J Matern Fetal Neonatal Med.* 2012 Oct;25 Suppl 3:49-52. doi: 10.3109/14767058.2012.712318. PMID: 23016618
- 37. GIANNI M, et al. The influence of a formula supplemented with dairy lipids and plant oils on the erythrocyte membrane omega-3 fatty acid profile in healthy full-term infants: A double-blind randomized controlled trial. *BMC Pediatr.* 2012 Oct 17;12:164. doi: 10.1186/1471-2431-12-164. PMID: 23072617; PMCID: PMC3480864
- 38. ROGGERO P, et al. Implementation of nutritional strategies decreases postnatal growth restriction in preterm infants. *PLoS ONE.* 2012, 7(12):e51166
- 39. BARTOK C. Babies fed breastmilk by breast versus by bottle: A pilot study evaluating early growth patterns. *Breastfeed Med.* 2011 Jun;6(3):117-24. doi: 10.1089/bfm.2010.0055. Epub 2010 Dec 6. PMID: 21133764 (2011)
- 40. ROGGERO P, et al. Rapid recovery of fat mass in small for gestational age preterm infants after term. *PLoS ONE.* 2011 Jan 5;6(1):e14489. doi: 10.1371/journal.pone.0014489. PMID: 21245927; PMCID: PMC301631
- 41. ANDERSON A, et al. Dietary trans fatty acid intake and maternal and infant adiposity. *Eur J Clin Nutr.* 2010 Nov;64(11):1308-15. doi: 10.1038/ejcn.2010.166. Epub 2010 Sep 8. PMID: 20823901
- 42. ANDERSON A. Association between infant feeding and early postpartum infant body composition: a pilot prospective study. *Int J Pediatr.* 2009;2009:648091. doi: 10.1155/2009/648091. Epub 2009 Mar 12. PMID: 20041019; PMCID: PMC2778447
- 43. ROGGERO P, et al. Influence of protein and energy intakes on body composition off formula-fed preterm infants after term [Short Communication]. *J Pediatr Gastroenterol Nutr.* 2008 Sep;47(3):375-8. doi: 10.1097/MPG.0b013e3181615cba. PMID: 18728539

\*\*\*\*\*