

## Validation

- FORSUM E, et al. An evaluation of the Pea Pod System for assessing body composition of moderately premature infants. *Nutrients*, 8(4):238 (2016)
- ROGGERO P, et al. Evaluation of air-displacement plethysmography for body composition assessment in preterm infants. *Pediatr Res*, 72(3):316-320 (2012)
- FRONDAS-CHAUTY A, et al. Air displacement plethysmography for determining body composition in neonates: validation using live piglets. *Pediatr Res*, 72(1):26-31 (2012)
- FIELDS D, et al. Body composition at 6 months of life: comparison of air displacement plethysmography and dual-energy x-ray Absorptiometry. *Obesity*, 20(11):2302-2306 (2012)
- ANDRES A, et al. Quantitative nuclear magnetic resonance to measure fat mass in infants and children. *Obesity*, 19(10):2089-2095 (2011)
- ELLIS K, et al. Body-composition assessment in infancy: air-displacement plethysmography compared with a reference 4-compartment model. *Am J Clin Nutr*, 79:653-660 (2007)
- YAO M, et al. Inter-device reliability of the PEA POD® for percent body fat estimates. <http://www.cosmed.com> (2005)
- MA G, et al. Validation of a new pediatric air-displacement plethysmograph for assessing Body Composition in Infants. *Am J Clin Nutr*, 79(4):653-660 (2004)
- URLANDO A, et al. A new air displacement plethysmograph for the measurement of body composition in infants. *Pediatr Res*, 53:486-492 (2003)
- SAINZ R, et al. Evaluation of a new pediatric air-displacement plethysmograph for body-composition assessment by means of chemical analysis of bovine tissue phantoms. *Am J Clin Nutr*, 77:364-370 (2003)
- YAO M, et al. Preliminary evaluation of a new pediatric air displacement plethysmograph for body composition assessment in infants. *Acta Diabetol*, 40:555-558 (2003)

## Validation of Other Techniques Using the PEA POD

- RODRIGUEZ-CANO AM, et al. Anthropometric and clinical correlates of fat mass in healthy term infants at 6 months of age. *BMC Pediatr*, 19(1):60 (2019)
- HAWKES C, et al. The relationship between IGF-I and -II concentrations and body composition at birth and over the first 2 months. *Pediatr Res*, 85(5): 687-692 (2019)
- JOSEFSON JL, et al. Fat mass estimation in neonates: Anthropometric models compared with air displacement plethysmography. *Br J Nutr*, 121(3):285-290 (2019)
- ROY SM, et al. Body Mass Index Is a Better Indicator of Body Composition than Weight-for-Length at Age 1 Month. *J Pediatr*, 204:77-83 e1 (2019)
- HUVANANDANA J, et al. An anthropometric approach to characterizing neonatal morbidity and body composition, using air displacement plethysmography as a criterion method. *PLoS One*, 12(3):e0195193 (2018)
- CHEN LW, et al. Which anthropometric measures best reflect neonatal adiposity? *Int J Obes*, 42(3):501-506 (2018)
- LIOTTO N, et al. Can basic characteristics estimate body composition in early infancy? *J Pediatr Gastroenterol Nutr*, 66(3):e76-e80 (2018)
- PERNG W, et al. An observational cohort study of weight- and length-derived anthropometric indicators with body composition at birth and 5 mo: The Healthy Start Study. *Am J Clin Nutr*, 106(2): 559-567 (2017)
- ROELANTS JA, et al. Foetal fractional thigh volume: an early 3D ultrasound marker of neonatal adiposity. *Pediatr Obes*, 12 (Suppl 1):65-71. (2017)

- CAUBLE JS, et al. Validity of anthropometric equations to estimate infant fat mass at birth and in early infancy. *BMC Pediatr*, 17(1):88 (2017)
- LARCADE J, et al. Estimation of fat-free mass at discharge in preterm infants fed with optimized feeding regimen. *J Pediatr Gastroenterol Nutr*, 64(1):115-118 (2017)
- RAMEL SE, et al. Do anthropometric measures accurately reflect body composition in preterm infants? *Pediatr Obes* Aug;12 (Suppl 1):72-77 (2017)
- HAWKES CP, et al. Body Composition within the first 3 months: optimized correction for length and correlation with BMI at 2 years. *Horm Res Paediatr*, 86(3):178-187 (2016)
- WROTTESELEY SV, et al. A comparison of body composition estimates using dual-energy X-ray absorptiometry and air-displacement plethysmography in South African neonates. *Eur J Clin Nutr*, 70(11):1254-1258 (2016)
- KIGER JR, et al. Preterm infant body composition cannot be accurately determined by weight and length. *J Neonatal Perinatal Med*, 9(3):285-90 (2016)
- RAUSCH I, et al. Reproducibility of MRI Dixon-based attenuation correction in combined PET/MR with applications for lean body mass estimation. *J Nucl Med*, 57(7):1096-101 (2016)
- TINT MT, et al. Estimation of fat-free mass in Asian neonates using bioelectrical impedance analysis. *Br J Nutr*, 115(6):1033-1042 (2016)
- BARBOUR LA, et al. Striking differences in estimates of infant adiposity by new and old DXA software, PEAPOD and skin-folds at 2 weeks and 1 year of life. *Pediatr Obes*, 11(4):264-71 (2016)
- MOORE GS, et al. Can Fetal Limb Soft Tissue Measurements in the Third Trimester Predict Neonatal Adiposity? *J Ultrasound Med*, 35(9):1915-24 (2016)
- WILBÆK R, et al. Calibration of bioelectrical impedance analysis for body composition assessment in Ethiopian infants using air-displacement plethysmography. *Eur J Clin Nutr*, 69(10):1099-104 (2015)
- DALY-WOLF KM, et al. Mid-arm circumference is a reliable method to estimate adiposity in preterm and term infants. *Pediatr Res*, 78(3):336-341 (2015)
- GRIJALVA-ETERNOD CS, et al. Midupper arm circumference and weight-for-length z scores have different associations with body composition: evidence from a cohort of Ethiopian infants. *Am J Clin Nutr*, 102(3):593-599 (2015)
- DE CUNTO A, et al. Can body mass index accurately predict adiposity in newborns? *Arch Dis Child Fetal Neonatal*, 99(3):F238-39 (2014)
- O'CONNOR C, et al. Fetal subcutaneous tissue measurements in pregnancy as a predictor of neonatal total body composition. *Prenatal Diag*, 34:1-4 (2014)
- ARIS I, et al. Body fat in Singaporean infants: development of body fat prediction equations in Asian newborns. *Eur J Clin Nutr*, 67(9):922-927 (2013)
- LINGWOOD B, et al. Prediction of fat-free mass and percentage of body fat in neonates using bioelectrical impedance analysis and anthropometric measures: validation against the PEA POD. *Br J Nutr*, 107(10):1545-1552 (2012)
- DEIERLEIN A, et al. An anthropometric model to estimate neonatal fat mass using air displacement plethysmography. *Nutr Metab*, 21:9-21 (2012)
- LEE W, et al. Fetal growth parameters and birth weight: their relationship to neonatal body composition. *Ultrasound Obstet Gynecol*, 33:441-446 (2009)

## Review

- BELFORT MB, et al. NICU Diet, Physical Growth and Nutrient Accretion, and Preterm Infant Brain Development. *Neoreviews*, 20(7):e385-e396 (2019)
- ANDREWS ET, et al. Measuring body composition in the preterm infant: Evidence base and practicalities. *Clin Nutr*, 38(6):2521-2530 (2019)

- STRYDOM K, et al. Factors affecting body composition in preterm infants: Assessment techniques and nutritional interventions. *Pediatr Neonatol*, 60(2):121-128 (2019)
- MAZAHERY H, et al. Air displacement plethysmography (pea pod) in full-term and pre-term infants: a comprehensive review of accuracy, reproducibility, and practical challenges. *Matern Health Neonatol Perinatol*, 4:12. doi: 10.1186/s40748-018-0079-z (2018)
- ROELANTS JA, et al. Prenatal markers of neonatal fat mass: a systematic review. *Clin Nutr*, 35(5):995-1007 (2016)
- TORO-RAMOS T, et al. Body composition during fetal development and infancy through the age of 5 years. *Eur J Clin Nutr*, 69(12):1279-1289 (2015)
- RICE et al. Neonatal body composition: measuring lean mass as a tool to guide nutrition management in the neonate. *Nutr Clin Pract*, 30(5):625-632 (2015)
- ZANINI RDE V, Body fat in children measured by DXA, air-displacement plethysmography, TBW and multicomponent models: a systematic review *Matern Child Health J*, 19(7):1567-1573 (2015)
- FIELDS DA, et al. Air displacement plethysmography: cradle to grave. *Nutr Clin Pract*, 30(2):219-226 (2015)
- HORAN M, et al. Methodologies to assess paediatric adiposity. *Ir J Med Sci*, 184(1):53-68 (2015)
- WELLS JCK, et al. Toward body composition reference data for infants, children, and adolescents. *Adv Nutr*, 5:320S-329S (2014)
- DEMERATH W, et al. Body composition assessment in the infant. *Am J Hum Biol*, 26(3):291-304 (2014)
- CORPELEIJN W, et al. Optimal growth of preterm infants. *World Rev Nutr Diet*, 106:149-155 (2013)
- WARD LC, et al. Assessing early growth and adiposity: report from an early nutrition academy workshop. *Ann Nutr Metab*, 63:120-130 (2013)
- LI C, et al. Infant body composition in the PEA POD® era: what have we learned and where do we go from here? *JDOHaD*, 4:116-120 (2013)
- WELLS J. Body composition in infants: evidence for developmental programming and techniques for measurement. *Rev Endocr Metab Disord*, 13:93-101 (2012)
- BARACOS V, et al. Advances in the science and application of body composition measurement. *JPEN*, 36(1):96-107 (2012)
- ELLIS K. Body composition in infancy: impact on health later in life. *Nestlé Nutr Inst Workshop Ser Pediatr Program*, 65:213-224 (2010)
- RUBIN L. Postnatal growth in preterm infants: too small, too big, or just right [Editorial]. *J Pediatr*, 54:473-475 (2009)
- ELLIS K. Evaluation of body composition in neonates and infants. *Semin Fetal Neonatal Med*, 12:87-91 (2007)
- ROGGERO P, et al. Measuring the body composition of preterm and term neonates: from research to clinical applications. *J Pediatr Gastroenterol Nutr*, 45:S159-162 (2007)
- WELLS J, et al. Programming of body composition by early growth and nutrition. *Pro Nutr Soc*, 66:423-434 (2007)
- RIGO J. Body composition during the first year of life. *Nestlé Nutr Inst Workshop Ser Pediatr Program*, 58:65-78 (2006)

### Normative Body Composition

- KURIYAN R, et al. The thin but fat phenotype is uncommon at birth in Indian babies. *J Nutr*, Dec 20. pii: nxz305. doi: 10.1093/jn/nxz305. [Epub ahead of print] (2019)
- WIECHERS C, et al. Neonatal body composition by air displacement plethysmography in healthy term singletons: a systematic review. *BMC Pediatr*, 19(1):489 (2019)
- WIECHERS C, et al. Neonatal body composition: cross-sectional study in healthy term singletons in Germany. *BMC Pediatr*, 19(1):488 (2019)
- ALEXANDER T, et al. Body composition of New Zealand-born term babies differs by ethnicity, gestational age and sex. *Early Hum Dev*, Nov 15;140:104924. doi: 10.1016/j.earlhumdev.2019.104924. [Epub ahead of print] (2019)

- NORRIS T, et al. New charts for the assessment of body composition, according to air-displacement plethysmography, at birth and across the first 6 mo of life. *Am J Clin Nutr*, 109(5): 1353-1360 (2019)
- FORSUM E, et al. Fat and fat-free mass of healthy Swedish children show tracking during early life, but there are differences. *Acta Paediatr*, 108(9):1704-1708 (2019)
- HENRICKSSON P, et al. Longitudinal assessment of body composition in healthy Swedish children from 1 week until 4 years of age. *Eur J Clin Nutr*, Aug 23. doi: 10.1038/ejcn.2017.125. [Epub ahead of print] (2017)
- VILLAR J, et al. Body composition at birth and its relationship with neonatal anthropometric ratios: the newborn body composition study of the INTERGROWTH-21st project. *Pediatr Res*, 82(2):305-316 (2017)
- SCHEURER JM, et al. Body composition trajectories from infancy to preschool in children born premature versus full-term. *J Pediatr Gastroenterol Nutr*, 64(6):e147-e153 (2017)
- DEMERATH EW, et al. New body composition reference charts for preterm infants. *Am J Clin Nutr*, 105(1):70-77 (2017)
- RAMEL S, et al. Body composition at birth in preterm infants between 30 and 36 weeks gestation. *Pediatr Obes*, 10(1):45-51 (2014)
- PAVIOTTI G, et al. Longitudinal growth and body composition of twins versus singleton in the first month of life. *Scientific World Journal*, Volume 2013, Article ID 108189, 3 pages (2013)
- CARBERRY A, et al. Is body fat percentage a better measure of undernutrition in newborns than birth weight percentiles? *Pediatr Res*, 74(6):730-736 (2013)
- ANDERSEN G, et al. Body composition from birth to 6 mo. of age in Ethiopian infants: reference data obtained by air-displacement plethysmography. *Am J Clin Nutr*, 98(4):885-894 (2013)
- LAW T, et al. Customized versus population-based growth curves: prediction of low body fat percent at term corrected gestational age following preterm birth. *J Matern Fetal Neonatal Med*, 25(7):1142-1147 (2012)
- FIELDS D, et al. Longitudinal body composition data in exclusively breast-fed infants: a multicenter study. *Obesity*, 19(9):1887-1891 (2011)
- HAWKES C, et al. Gender-and gestational age-specific body fat percentage at birth. *Pediatrics*, 128(3):e645-651 (2011)
- JOHNSON M, et al. Preterm birth and body composition at term equivalent age: a systematic review and meta-analysis. *Pediatrics*, 130(3):e640-649 (2011)
- ROGGERO P, et al. Quality of growth in exclusively breast-fed infants in the first six months of life: an Italian study. *Pediatr Res*, 68:542-544 (2010)
- CARBERRY A, et al. Body composition from birth to 4.5 months in infants born to non-obese women. *Pediatr Res*, 68(1):84-88 (2010)

### **Body Composition Observation in Preterm Infants**

- CHMIELEWSKA A, et al. Lean tissue deficit in preterm infants persists up to 4 months of age: results from a Swedish longitudinal study. *Neonatology*, Dec 10:1-8. doi: 10.1159/000503292. [Epub ahead of print] (2019)
- FORSUM EK, et al. Premature birth was not associated with increased body fatness in four-year-old boys and girls. *Acta Paediatr*, Aug 28. doi: 10.1111/apa.14990. [Epub ahead of print] (2019)
- BELL KA et al. Associations of growth and body composition brain size in preterm infants. *J Pediatr*. Jul 31. pii: S0022-3476(19)30826-1. doi: 10.1016/j.jpeds.2019.06.062. [Epub ahead of print] (2019)
- MC CLUNAN K, et al. Effect of nutritional intake on the body composition of HIV-exposed and HIV-unexposed preterm and low birth weight infants. *Breastfeed Med*, 14(3):144-153 (2019)
- JEROME M, et al. Racial disparities in body composition of preterm infants. *Curr Dev Nutr*, Jun 13;3(Suppl 1). pii: nzz041.P21-049-19. doi: 10.1093/cdn/nzz041.P21-049-19 (2019)

- MORNIROLI D, et al. Is the body composition development in premature infants associated with a distinctive nuclear magnetic resonance metabolomic profiling of urine? *J Matern Fetal Neonatal Med*, 32(14):2310-2318 (2019)
- DISSANAYAKE HU, et al. Body Fatness and Cardiovascular Health in Newborn Infants, *J Clin Med*, Sep 11;7(9). Pii: E270. Doi: 10.3390/jcm7090270 (2018)
- PIEMONTESE P, et al. The Effect of Human Milk on Modulating the Quality of Growth in Preterm Infants. *Front Pediatr*, Oct 9;6:291. doi: 20.3389/fped.2018.00291. eCollection (2018)
- SCHEURER JM, et al. Body Composition Changes from Infancy to 4 Years and Associations with Early Childhood Cognition in Preterm and Full-Term Children. *Neonatology*, 114(2):169-176 (2018)
- DA SILVA MARTINS A, et al. Growth and body composition in preterm newborns with bronchopulmonary dysplasia: a cohort study. *J Perinat Med*, 46(8):913-918 (2018)
- VILLELA LD, et al. Body composition in preterm infants with intrauterine growth restriction: A cohort study. *J Perinat Med*, 46(7):804-810 (2018)
- VILLELA LD, et al. Growth and body composition of preterm infants less than or equal to 32 weeks: Cohort study. *Early Hum Dev*, 117:90-95 (2018)
- ALGOTAR A, et al. Unique patterns of body composition and anthropometric measurements during maturation in neonatal intensive care unit neonates: Opportunities for modifying nutritional therapy and influencing clinical outcomes. *J Parenter Enteral Nutr*, 42(1):231-238 (2018)
- ABERA M, et al. Relation between body composition at birth and child development at 2 years of age: a prospective cohort study among Ethiopian children. *Eur J Clin Nutr*, 71(12):1411-1417 (2017)
- PAVIOTTI G, et al. Higher growth, fat and fat-free masses correlate with larger cerebellar volumes in preterm infants at term. *Acta Paediatr*, 106(6):918-925 (2017)
- PIYASENA C, et al. Dynamic Changes in DNA Methylation Occur during the First Year of Life in Preterm Infants, *Front Endocrinol (Lausanne)*, 15;7:158 (2016)
- RAMEL SE, et al. Greater early gains in fat-free mass, but not fat mass, are associated with improved neurodevelopment at 1 year corrected age for prematurity in very low birth weight preterm infants. *J Pediatr*, 173:108-115 (2016)
- SCHEURER JM, et al. Diminished growth and lower adiposity in hyperglycemic very low birth weight neonates at 4 months corrected age. *J Perinatol*, 36(2):145-150 (2016)
- GIANNI ML, et al. Body composition in late preterm infants according to percentile at birth. *Pediatr Res*, 79(5):710-715 (2016)
- DE CUNTO, et al. Impact of surgery for neonatal gastrointestinal diseases on weight and fat mass. *J Pediatr*, 167(3):568-571 (2015)
- ROGGERO P, et al. No relative increase in intra-abdominal adipose tissue in healthy unstressed preterm infants at term. *Neonatology*, 107(1):14-19 (2015)
- PIYASENA C, et al. Dynamics of DNA methylation at IGF2 in preterm and term infants during the first year of life: an observational study. *Lancet*, 26;385 Suppl 1:S81 (2015)
- OHLAGER E, et al. Body composition in late preterm infants in the first 10 days of life and at full term. *Acta Paediatr*, 103(7):737-743 (2014)
- SIMON L, et al. Determinants of body composition in preterm infants at the time of hospital discharge. *Am J Clin Nutr*, 100:98-104 (2014)
- MEYERS J, et al. Potential influence of total parenteral nutrition on body composition at discharge in preterm infants. *J Matern Fetal Neonatal Med*, 26(15):1548-1553 (2013)
- PFISTER K, et al. Exploratory study of the relationship of fat-free mass to speed of brain processing in preterm infants. *Pediatr Res*, 74(5):576-583 (2013)
- SIMON L, et al. Effect of sex and gestational age on neonatal body composition. *Br J Nutr*, 109:1105-1108 (2013)



- LIOTTO N, et al. Growth and body composition changes in late preterm infants in the first months of life. *Pediatr Med Chir*, 35(4):172-176 (2013)
- PIEMONTESE P, et al. Effect of prematurity on fat mass distribution and blood pressure at prepubertal age: a follow-up study. *Pediatr Med Chir*, 35(4):166-171 (2013)
- BARANOV A, et al. Evaluation of premature infants' nutritional status by air plethysmography: first Russian prospective study. *Vestn Ross Akad Med Nauk*, (4)10-16 (2013)
- TARONI F, et al. Intra-abdominal adiposity in preterm infants: an explorative study. *Pediatr Med Chir*, 34(6):283-286 (2012)
- GIANNI L, et al. Postnatal catch-up fat after late preterm birth. *Pediatr Res*, 72(6):637-640 (2012)
- LAW T, et al. Ultrasound assessment of intrauterine growth restriction: relationship to neonatal body composition. *Am J Obstet Gynecol*, 205(3):255.e1-6 (2011)
- RAMEL S, et al. Body composition changes in preterm infants following hospital discharge: comparison with term infants. *J Pediatr Gastroenterol Nutr*, 53(3):333-338 (2011)
- ROGGERO P, et al. Is term newborn body composition being achieved postnatally in preterm infants? *Early Hum Dev*, 85(6):349-352 (2009)
- GIANNI M, et al. Adiposity in small for gestational age preterm infants assessed at term equivalent age. *Arch Dis Child Fetal Neonatal Ed*, 94(5):F368-372 (2009)
- ROGGERO P, et al. Postnatal growth failure in preterm infants: recovery of growth and body composition after term. *Early Hum Dev*, 84(8):555-559 (2008)
- TARONI F, et al. Body composition in small for gestational age newborns. *Pediatr Med Chir*, 30(6):296-301 (2008)

### **Body Composition Observation in Term Infants**

- KADAKIA R, et al. Association of cord blood methylation with neonatal leptin: an epigenome wide association study. *PLoS One*, 14(12): e0226555 (2019)
- DAVIS SM, et al. Sex differences in infant body composition emerge in the first 5 months of life. *J Pediatr Endocrinol Metab*, 32(11):1235-1239 (2019)
- RODRÍGUEZ-CANO AM, et al. Higher fat mass and fat mass accretion during the first six months of life in exclusively breastfed infants. *Pediatr Res*, Aug 21. doi: 10.1038/s41390-019-0542-1. [Epub ahead of print] (2019)
- WIBAEK R, et al. Associations of fat mass and fat-free mass accretion in infancy with body composition and cardometabolic risk markers at 5 years: The Ethiopian iABC birth cohort study. *PLoS Med*, 20;16(8):e1002888 (2019)
- SMITH HA, et al. Difference between body composition of formula-and breastfed infants at birth. *J Dev Orig Health Dis*, May 28:1-5 doi: 10.1017/S2040174419000187. [Epub ahead of print] (2019)
- LARSSON A, et al. Body composition and growth in full-term small for gestational age and large for gestational age Swedish infants assessed with air displacement plethysmography at birth and at 3-4 months of age. *PLoS One*, 14(5):e0207978 (2019)
- KADAKIA R, et al. Cord Blood Metabolites Associated with Newborn Adiposity and Hyperinsulinemia. *J Pediatr*, 203:144-149 e1 (2018)
- EUCLYDES VLV, et al. Cord blood concentrations of leptin, zinc- $\alpha$ 2-glycoprotein, and adiponectin, and adiposity gain during the first 3 mo of life. *Nutrition*, 54:89-93 (2018)
- ANDERSEN GS, et al. Body Composition Growth Patterns in Early Infancy: A Latent Class Trajectory Analysis of the Ethiopian iABC Birth Cohort. *Obesity*, 26(7):1225-1233 (2018)
- ADMASSU B, et al. Body composition during early infancy and its relation with body composition at 4 years of age in Jimma, an Ethiopian prospective cohort study. *Nutr Diabetes*, 8(1):46 (2018)

- ABERA M, et al. Body composition during early infancy and developmental progression from 1 to 5 years of age: The Infant Anthropometry and Body Composition (iABC) cohort study among Ethiopian children. *Br J Nutr*, 119(11):1263-1273 (2018)
- DISSANAYAKE HU, et al. Noninvasive assessment of autonomic function in human neonates born at the extremes of fetal growth spectrum. *Physiol Rep*, 6(8):e13682 (2018)
- ADMASSU B, et al. Accretion of Fat-Free Mass Rather Than Fat Mass in Infancy Is Positively Associated with Linear Growth in Childhood. *J Nutr*, 148(4):607-615 (2018)
- SCHNEIDER CR, et al. Associations of neonatal adiponectin and leptin with growth and body composition in African American infants. *Pediatr Obes*, 13(8):485-491 (2018)
- KADAKIA R, et al. Maternal pre-pregnancy BMI downregulates neonatal cord blood LEP methylation. *Pediatr Obes*, S1:57-64 (2017)
- ADMASSU B, et al. Body composition at birth and height at 2 years: a prospective cohort study among children in Jimma, Ethiopia. *Pediatr Res*, 82(2):209-214 (2017)
- SAUDER KA, et al. Predictors of Infant Body Composition at 5 Months of Age: The Healthy Start Study. *J Pediatr*, 183:94-99.e1 (2017)
- BREIJ LM, et al. Appetite-regulating hormones in early life and relationships with type of feeding and body composition in healthy term infants. *Eur J Nutr*, 56(4):1725-1732 (2017)
- BREIJ LM, et al. Longitudinal fat mass and visceral fat during the first 6 months after birth in healthy infants: support for a critical window for adiposity in early life. *Pediatr Obes*, 12(4):286-294 (2017)
- CHAOIMH CN, et al. Cord blood leptin and gains in body weight and fat mass during infancy. *Eur J Endocrinol*, Nov;175(5):403-10 (2016)
- KADAKIA R, et al. Neonatal adiposity increases with rising cord blood IGF-1 levels. *Clin Endocrinol (Oxf)*, 85(1):70-5 (2016)
- O'DONOVAN SM, et al. Neonatal adiposity increases the risk of atopic dermatitis during the first year of life. *J Allergy Clin Immunol*, 137(1):108-117 (2016)
- PALEY C, et al. Body fat differences by self-reported race/ethnicity in healthy term newborns. *Pediatr Obes*, 11(5):361-8 (2016)
- FAROOQ M, et al. Monitoring of infant feeding behavior using a jaw motion sensor. *J Healthc Eng*, 6(1):23-40 (2015)
- O'DONOVAN SM, et al. Cohort profile: The Cork BASELINE birth cohort study: babies after SCOPE: evaluating the longitudinal impact on neurological and nutritional endpoints. *Int J Epidemiol* 44(3):764-775 (2015)
- HENRICKSSON P, et al. Variation in the fat mass and obesity-related (FTO) genotype is not associated with body fatness in infants, but possibly with their length. *Ped Obes*, 9(5):e112-115 (2014)
- DONNELLEY EL, et al. Antenatal predictors and body composition of large-for-gestational-age newborns: perinatal health outcomes. *J Perinatol*, 34(9):698-704 (2014)
- O'CONNOR C, et al. Birth weight and neonatal adiposity prediction using fractional limb volume obtained with 3D ultrasound. *Fetal Diag Ther*, 36(1):44-48 (2014)
- GIANNI M, et al. Body composition changes in the first 6 months of life according to method of feeding. *J Hum Lact*, 30(2):148-155 (2014)
- ERIKSSON B, et al. Body-composition development during early childhood and energy expenditure in response to physical activity in 1.5-year-old children. *Am J Clin Nutr*, 96(3):567-73 (2012)
- STANFIELD K, et al. Differences in body composition between infants of South Asian and European ancestry: the London Mother and Baby Study. *Int J Epidemiol*, 41(5):1409-1418 (2012)
- LEE W, et al. The relationship of newborn adiposity to fetal growth outcome based on birth weight or the modified neonatal growth assessment score. *J Matern Fetal Neonatal Med*, 25(10):1933-1940 (2012)
- ANDERSEN G, et al. Fat and fat-free mass at birth: air displacement plethysmography measurements on 350 Ethiopian newborns. *Pediatr Res*, 70:501-506 (2011)

- ERIKSSON B, et al. Fat-free mass hydration in newborns: assessment and implications for body composition studies. *Acta Paediatr*, 100:680-686 (2011)
- ROGGERO P, et al. Neonatal period: body composition changes in breast-fed full-term newborns. *Neonatology*, 97:139-143 (2010)
- ERIKSSON B, et al. Body composition in full-term healthy infants measured with air displacement plethysmography at 1 and 12 weeks of age. *Acta Paediatr*, 99(4):563-568 (2009)
- FIELDS D, et al. Sex differences in body composition early in life. *Gender Med*, 6(2):369-375 (2009)
- MOYER-MILEUR L, et al. Newborn adiposity measured by plethysmography is not predicted by late gestation two-dimensional ultrasound measures of fetal growth. *J Nutr*, 139:1772-1778 (2009)

### Mother/Newborn Relationship

- DIAZ EC, et al. Parental adiposity differentially associates with newborn body composition. *Pediatr Obes*, Dec 19:e12596. doi: 10.1111/ijpo.12596. [Epub ahead of print] (2019)
- NEHAB SRG, et al. Influence of gestational and perinatal factors on body composition of full-term newborns. *J Pediatr (Rio J)*, Nov 9. pii: S0021-7557(19)30391-2. doi: 10.1016/j.jpmed.2019.09.006. [Epub ahead of print] (2019)
- BENNETT AE, et al. Maternal sociodemographic and health behaviours associated with adiposity in infants as measured by air displacement plethysmography. *Early Hum Dev*, Oct 26;140:104887. doi: 10.1016/j.earlhumdev.2019.104887. [Epub ahead of print] (2019)
- HOLLANDERS J, et al. No association between glucocorticoid diurnal rhythm in breastmilk and infant body composition at 3 months. *Nutrients*, Oct 2;11(10). pii: E2351. doi: 10.3390/nu11102351 (2019)
- ABREU LRS, et al. Gestational diabetes mellitus, pre-pregnancy body mass index, and gestational weight gain as risk factors for increased fat mass in Brazilian newborns. *PLoS One*, 14(8):e0221971 (2019)
- MUDD LM, et al. Relations among maternal physical activity during pregnancy and child body composition. *Obes Sci Prac*, 5(3):246-250 (2019)
- FUENTES ED, et al. Parental obesity differentially associates with newborn adiposity. Is programming of the GH-IGF axis involved? *Curr Dev Nutr*, Jun 13;3(Suppl 1). pii: nzz041.OR09-05-19. doi: 10.1093/cdn/nzz041.OR09-05-19 (2019)
- VAN LEE L, et al. Prospective associations of maternal choline status with offspring body composition in the first 5 years of life in two large mother-offspring cohorts: the Southampton Women's Survey cohort and the Growing Up in Singapore Towards healthy Outcomes cohort. *Int J Epidemiol*, 48(2):433-444 (2019)
- ANDERSSON-HALL UK, et al. Maternal obesity and gestational diabetes mellitus affect body composition through infancy: the PONCH study. *Pediatr Res*, 85(3):369-377 (2019)
- PHANG M, et al. Increased  $\alpha$ -linolenic acid intake during pregnancy is associated with higher offspring birth weight. *Curr Dev Nutr*, 2018 Oct 18;3(2):nzy081. doi: 10.1093/cdn/nzy081. eCollection Feb (2019)
- MOORE BF, et al. Fetal exposure to maternal active and secondhand smoking with offspring early-life growth in the Healthy Start study. *Int J Obes*, 43(4):652-662 (2019)
- KENNEDY RAK, et al. The relationship between early pregnancy dietary intakes and subsequent birthweight and neonatal adiposity. *J Pub Health*, 40(4):747-755 (2018).
- WROTTESELEY SV, et al. Maternal traditional dietary pattern and antiretroviral treatment exposure are associated with neonatal size and adiposity in urban, black South Africans. *Br J Nutr*, 120(5):557-566 (2018)
- BARBOUR LA, et al. Postprandial Triglycerides Predict Newborn Fat More Strongly than Glucose in Women with Obesity in Early Pregnancy. *Obesity*, 26(8):1347-1356 (2018)



- MACAULAY S, et al. The effects of gestational diabetes mellitus on fetal growth and neonatal birth measures in an African cohort. *Diabet Med*, 35(10):1425-1433 (2018)
- MOORE BF, et al. Proinflammatory diets during pregnancy and neonatal adiposity in the Healthy Start Study. *J Pediatr*, 195:121-127 (2018)
- DAHLY DL, et al. Associations between maternal lifestyle factors and neonatal body composition in the Screening for Pregnancy Endpoints (Cork) cohort study. *Int J Epidemiol*, 47(1):131-145 (2018)
- GALLAGHER D, et al. Greater neonatal fat-free mass and similar fat mass following a randomized trial to control excess gestational weight gain. *Obesity*, 26(3):578-587 (2018)
- CHIA AR, et al. Adherence to a healthy eating index for pregnant women is associated with lower neonatal adiposity in a multiethnic Asian cohort: The Growing Up in Singapore Towards Healthy Outcomes (GUSTO) Study. *Am J Clin Nutr*, 107(1):71-79 (2018)
- BAKER PR 2<sup>nd</sup>, et al. Maternal obesity and increased neonatal adiposity correspond with altered infant mesenchymal stem cell metabolism. *JCI Insight*, Nov 2. 2(21). pii: 94200. doi: 10.1172/jci.insight.94200. [Epub ahead of print] (2017)
- DAHLY DL, et al. Associations between maternal lifestyle factors and neonatal body composition in the Screening for Pregnancy Endpoints (Cork) cohort study. *Int J Epidemiol*, Nov 10. doi: 10.1093/ije/dyx221. [Epub ahead of print] (2017)
- MOORE BF, et al. Proinflammatory Diets during Pregnancy and Neonatal Adiposity in the Healthy Start Study. *J Pediatr*. Dec 5. pii: S0022-3476(17)31379-3. doi: 10.1016/j.jpeds.2017.10.030. [Epub ahead of print] (2017)
- BAKER PR 2<sup>nd</sup>, et al. Altered gene expression and metabolism in fetal umbilical cord mesenchymal stem cells correspond with differences in 5-month-old infant adiposity gain. *Sci Rep*, 7(1):18095 (2017)
- BREIJ LM, et al. Impact of Early Infant Growth, Duration of Breastfeeding and Maternal Factors on Total Body Fat Mass and Visceral Fat at 3 and 6 Months of Age. *Ann Nutr Metab*, 71(3-4):203-210 (2017)
- NORRIS T, et al. Do changing levels of maternal exercise during pregnancy affect neonatal adiposity? Secondary analysis of the babies after SCOPE: evaluating the longitudinal impact using neurological and nutritional endpoints (BASELINE) birth cohort (Cork, Ireland). *BMJ Open*, 7(11):e017987 (2017)
- MCKENZIE KM, et al. Quantity and Quality of Carbohydrate Intake during Pregnancy, Newborn Body Fatness and Cardiac Autonomic Control: Conferred Cardiovascular Risk? *Nutrients*, 9(12) (2017)
- MACDONALD TM, et al. Reduced growth velocity across the third trimester is associated with placental insufficiency in fetuses born at a normal birthweight: a prospective cohort study. *BMC Med*, 15(1):164 (2017)
- RUDOLPH MC, et al. Early infant adipose deposition is positively associated with the n-6 to n-3 fatty acid ratio in human milk independent of maternal BMI. *Int J Obes (Lond)*, 41(4):510-517 (2017)
- BERNARD JY, et al. Maternal plasma phosphatidylcholine polyunsaturated fatty acids during pregnancy and offspring growth and adiposity. *Prostaglandins Leukot Essent Fatty Acids*, 121:21-29 (2017)
- STARLING AP, et al. Perfluoroalkyl substances during pregnancy and offspring weight and adiposity at birth: examining mediation by maternal fasting glucose in the Healthy Start Study. *Environ Health Perspect*, Jun 26;125(6):067016 (2017)
- SAUDER KA, et al. Prenatal vitamin D intake, cord blood 25-hydroxyvitamin D, and offspring body composition: The Healthy Start Study. *Nutrients*, Jul 22;9(7) (2017)
- FUGLESTAD AJ, et al. Maternal executive function, infant feeding responsiveness and infant growth during the first 3 months. *Pediatr Obes*, Aug;12 Suppl 1:102-110 (2017)
- MOORE BF, et al. Exposure to secondhand smoke, exclusive breastfeeding and infant adiposity at age 5 months in the Healthy Start study. *Pediatr Obes*, Aug;12 Suppl 1:111-119 (2017)
- CASTRO NP, et al. The relationship between maternal plasma leptin and adiponectin concentrations and newborn adiposity. *Nutrients*, 23;9(3) (2017)

- LOGAN KM, et al. Diabetes in pregnancy and infant adiposity: systematic review and meta-analysis. *Arch Dis Child Fetal Neonatal Ed*, 102(1):F65-F72 (2017)
- PEREIRA-DA-SILVA L, et al. Low adherence to Mediterranean diet in Portugal: pregnant women nutrition in Portugal and its repercussions. *Acta Med Port*, 29(10):658-666 (2016)
- OKESENE-GAFA K, et al. A randomised controlled demonstration trial of multifaceted nutritional intervention and or probiotics: the healthy mums and babies (HUMBA) trial. *BMC Pregnancy Childbirth*, 16(1):373 (2016)
- SIMONI J, et al. A comparison of intestinal microbiota in a population of low-risk infants exposed and not exposed to intrapartum antibiotics: The Baby & Microbiota of the Intestine cohort study protocol. *BMC Pediatr*, 16(1):183 (2016)
- JOSEFSON JL, et al. Excessive gestational weight gain in the first trimester among women with normal glucose tolerance and resulting neonatal adiposity. *J Perinatol*, 36(12):1034-1038 (2016)
- CRUME TL, et al. Maternal dietary intake during pregnancy and offspring body composition: The Healthy Start Study. *Am J Obstet Gynecol*, 215(5):609.e1-609.e8 (2016)
- SHAPIRO AL, et al. Nicotinamide promotes adipogenesis in umbilical cord-derived mesenchymal stem cells and is associated with neonatal adiposity: The Healthy Start BabyBUMP Project. *PLoS One*, 11(7):e0159575 (2016)
- SHAPIRO AL, et al. Maternal diet quality in pregnancy and neonatal adiposity: The Healthy Start Study. *Int J Obes*, 40(7):1056-62 (2016)
- LEMAS DJ, et al. Alterations in human milk leptin and insulin are associated with early changes in the infant intestinal microbiome. *Am J Clin Nutr*, 103(5):1291-1300 (2016)
- KIZIRIAN NV, et al. Macronutrient balance and dietary glycemic index in pregnancy predict neonatal body composition. *Nutrients*, 8(5):E270 (2016)
- MCCARTHY FP, et al. Parental physical and lifestyle factors and their association with newborn body composition. *BJOG*, 123(11):1824-9 (2016)
- KIZIRIAN NV, et al. Effects of a low-glycemic index diet during pregnancy on offspring growth, body composition, and vascular health: a pilot randomized controlled trial. *Am J Clin Nutr*, 103(4):1073-1082 (2016)
- TINIUUS RA, et al. Maternal inflammation during late pregnancy is lower in physically active compared with inactive obese women. *Appl Physiol Nutr Metab*, 41(2):191-198 (2016)
- MARKOVIC TP, et al. Randomized controlled trial investigating the effects of a low-glycemic index diet on pregnancy outcomes in women at high risk of gestational diabetes mellitus: the GI Baby 3 Study. *Diabetes Care*, 39(1):31-38 (2016)
- SAUDER KA, et al. Exploring the association between maternal prenatal multivitamin use and early infant growth: The Healthy Start Study. *Pediatr Obes*, 11(5):434-41 (2016)
- HULL HR, et al. Maternal obesity influences the relationship between location of neonate fat mass and total fat mass. *Pediatr Obes*, 10(4):245-51 (2015)
- TINIUUS RA, et al. Altered maternal lipid metabolism is associated with higher inflammation in obese women during late pregnancy. *Integr Obes Diabetes*, 2(1):168-175 (2015)
- BREIJ LM, et al. Maternal and fetal determinants of neonatal body composition. *Horm Res Paediatr*, 84(6):388-395 (2015)
- LEMAS DJ, et al. Associations of maternal weight status prior and during pregnancy with neonatal cardiometabolic markers at birth: The Healthy Start Study. *Int J Obes (Lond)*, 39(10):1437-1442 (2015)
- HENRIKSSON P, et al. Gestational weight gain according to Institute of Medicine recommendations in relation to infant size and body composition. *Pediatr Obes*, 10(5):388-394 (2015)
- HENRIKSSON P, et al. Glucose homeostasis variables in pregnancy versus maternal and infant body composition. *Nutrients*, 7(7):5615-5627 (2015)

- HENRIKSSON P, et al. Parental fat-free mass is related to the fat-free mass of infants and maternal fat mass is related to the fat mass of infant girls. *Acta Paediatr*, 104(5):491-497 (2015)
- STARLING AP, et al. Associations of maternal BMI and gestational weight gain with neonatal adiposity in The Healthy Start Study. *Am J Clin Nutr*, 101(2):302-309 (2015)
- HARROD CS, et al. Exposure to prenatal smoking and early-life body composition: The Healthy Start Study. *Obesity*, 23(1):234-241 (2015)
- CRUME TL, et al. Maternal fuels and metabolic measures during pregnancy and neonatal body composition: The Healthy Start Study. *J Clin Endocrinol Metab*, 100(4):1672-1680 (2015)
- HARROD CS, et al. Quantity and timing of maternal prenatal smoking on neonatal body composition: the Healthy Start Study. *J Pediatr*, 165(4):707-712 (2014)
- HARROD CS, et al. Physical activity in pregnancy and neonatal body composition: The Healthy Start Study. *Obstet Gynecol*, 124(2 Pt 1):257-264 (2014)
- ESTAMPADOR A, et al. Infant body composition and adipokine concentrations in relation to maternal gestational weight gain. *Diabetes Care*, 37(5):1432-1438 (2014)
- PEREIRA-DA-SILVA L, et al. The adjusted effect of body mass index, energy and macronutrient intakes during pregnancy, and gestational weight gain on body composition of full-term neonates. *Am J Perinatol*, 31(10):875-882 (2014)
- JOSEFSON JL, et al. Maternal obesity and vitamin D sufficiency are associated with cord blood vitamin D insufficiency. *J Clin Endocrinol Metab*, 98(1):114-9 (2013)
- JOSEFSON JL, et al. Maternal leptin predicts adiposity of the neonate. *Horm Res Paediatr*, 81:13-19 (2013)
- POMEROY J, et al. Maternal physical activity and insulin action in pregnancy and their relationships with infant body composition. *Diabetes Care*, 36:267-269 (2013)
- AU C, et al. Body composition is normal in term infants born to mothers with well-controlled gestational diabetes mellitus. *Diabetes Care*, 36(9):e164 (2013)
- CARBERRY A, et al. Customized versus population-based birth weight charts for the detection of neonatal growth and perinatal morbidity in a cross-sectional study of term neonates. *Am J Epidemiol*, 178(8):1301-1308 (2013)
- JOSEFSON J, et al. Excessive weight gain in women with a normal pre-pregnancy BMI is associated with increased neonatal adiposity. *Ped Obes*, 8(2):e33-36 (2013)
- AU C, et al. Fetal and maternal factors associated with neonatal adiposity as measured by air displacement plethysmography: a large cross-sectional study. *Early Hum Dev*, 89(10):839-843 (2013)
- ANDRES A, et al. Body fat mass of exclusively breastfed infants born to overweight mothers. *J Acad Nutr Diet*, 112:991-995 (2012)
- LARSON ODE K, et al. Decelerated early growth in infants of overweight and obese mothers. *J Pediatr*, 161(6):1028-1034 (2011)
- HULL H, et al. Higher infant body fat with excessive gestational weight gain in overweight women. *Am J Obstet Gynecol*, 205(3):211.e1-7 (2011)
- HULL H, et al. Impact of maternal body mass index on neonate birthweight and body composition. *Am J Obstet Gynecol*, 198(4):416.e1-6 (2008)

### **Body Composition and Nutritional Intervention**

- ATCHLEY CB, et al. Enhanced protein diet for preterm infants: A prospective, randomized, double-blind, controlled trial. *J Pediatr Gastroenterol Nutr*, 69(2):218-22 (2019)
- MACEDO I, et al. Associations of measured protein and energy intakes with growth and adiposity in human milk-fed preterm infants at term postmenstrual age: A cohort study. *Am J Perinatol*, 35(9):882-891 (2018)

- MORLACCHI L, et al. Protein use and weight-gain quality in very-low-birth-weight preterm infants fed human milk or formula. *Am J Clin Nutr*, 107(2):195-200 (2018)
- LIOTTO N, Clinical evaluation of two different protein content formulas fed to full-term healthy infants: a randomized controlled trial. *BMC Pediatr*, 18(1): 59 (2018)
- TELLER IC, et al. Post-discharge formula feeding in preterm infants: a systematic review mapping evidence about the role of macronutrient enrichment. *Clin Nutr*, 35(4):791-801 (2016)
- GIANNI ML, et al. Does human milk modulate body composition in late preterm infants at term-corrected age? *Nutrients*, 8(10):664 (2016)
- BRENNAN AM, et al. Optimising preterm nutrition: present and future. *Proc Nutr Soc*, 75(2):154-161 (2016)
- MCCLEOD G, et al. Comparing different methods of human breast milk fortification using measured v. assumed macronutrient composition to target reference growth: a randomised controlled trial. *Br J Nutr*, 115(3):431-439 (2016)
- BLOOMFIELD FH, et al. The ProVIDe study: the impact of protein intravenous nutrition on development in extremely low birthweight babies. *BMC Pediatr*, 26;15:100 (2015)
- PEREIRA-DA-SILVA L, et al. Resting energy expenditure, macronutrient utilization, and body composition in term infants after corrective surgery of major congenital anomalies: A case-study. *J Neonatal Perinatal Med*, 8(4):403-412 (2015)
- MCLEOD G, et al. Feasibility study: assessing the influence of macronutrient intakes on preterm body composition, using air displacement plethysmography. *J Paediatr Child Health*, 51(9):862-869 (2015)
- PEREIRA-DA-SILVA L, et al. The effect of long-chain polyunsaturated fatty acids intake during pregnancy on adiposity of healthy full-term offspring at birth. *J Perinatol*, 35(3):177-180 (2015)
- GIANNI ML, et al. Formula-fed infants have significantly higher fat-free mass content in their bodies than breastfed babies. *Acta Paediatr*, 103(7):e277-281 (2014)
- TIMBY N, et al. Neurodevelopment, nutrition, and growth until 12 mo. of age in infants fed a low-energy, low-protein formula supplemented with bovine milk fat globule membranes: a randomized controlled trial. *Am J Clin Nutr*, 99:860-868 (2014)
- GIANNI ML, et al. The influence of a formula supplemented with dairy lipids and plant oils on the erythrocyte membrane omega-3 fatty acid profile in healthy full-term infants: a double-blind randomized controlled trial. *BMC Pediatr*, 12:164 (2012)
- ROGGERO P, et al. Implementation of nutritional strategies decreases postnatal growth restriction in preterm infants. *PLoS ONE*, 7(12):e51166 (2012)
- ROGGERO P, et al. Growth and fat-free mass gain in preterm infants after discharge: a randomized controlled trial. *Pediatrics*, 130:e1215-1221 (2012)
- ROGGERO P, et al. Effect of nutrition on growth and body composition in infants born preterm. *J Matern Fetal Neonatal Med*, 25(53):49-52 (2012)
- ROGGERO P, et al. Rapid recovery of fat mass in small for gestational age preterm infants after term. *PLoS ONE*, 6(1):e14489 (2011)
- BARTOK C. Babies fed breastmilk by breast versus by bottle: a pilot study evaluating early growth patterns. *Breastfeed Med*, 6(3):117-124 (2011)
- ANDERSON A, et al. Dietary trans fatty acid intake and maternal and infant adiposity. *Eur J Clin Nutr*, 64:1308-1315 (2010)
- ANDERSON A. Association between infant feeding and early postpartum infant body composition: a pilot prospective study. *Int J Pediatr*, Volume 2009, Article ID 648091, 7 pages (2009)
- ROGGERO P, et al. Influence of protein and energy intakes on body composition off formula-fed preterm infants after term [Short Communication]. *J Pediatr Gastroenterol Nutr*, 47(3):375-378 (2008)